

On representations of Higher Spin symmetry algebras for mixed-symmetry HS fields on AdS-spaces. Lagrangian formulation

Č. Burdík^a and A. Reshetnyak^b

^aDepartment of Mathematics, Czech Technical University, Prague 12000, Czech Republic,

^bInstitute of Strength Physics and Materials Science, Tomsk 634021, Russia

E-mail: ^aburdik@kmlinux.fjfi.cvut.cz, ^breshet@ispms.tsc.ru

Abstract. We derive non-linear commutator HS symmetry algebra, which encode unitary irreducible representations of AdS group subject to Young tableaux $Y(s_1, \dots, s_k)$ with $k \geq 2$ rows on d -dimensional anti-de-Sitter space. Auxiliary representations for specially deformed non-linear HS symmetry algebra in terms of generalized Verma module in order to additively convert a subsystem of second-class constraints in the HS symmetry algebra into one with first-class constraints are found explicitly for the case of HS fields for $k = 2$ Young tableaux. The oscillator realization over Heisenberg algebra for obtained Verma module is constructed. The results generalize the method of auxiliary representations construction for symplectic $sp(2k)$ algebra used for mixed-symmetry HS fields on a flat spaces and can be extended on a case of arbitrary HS fields in AdS-space. Gauge-invariant unconstrained reducible Lagrangian formulation for free bosonic HS fields with generalized spin (s_1, s_2) is derived.

1. Introduction

Increased interest to higher-spin field theory is mainly conditioned by expected output of LHC on the planned capacity. It suspects not only the proof of supersymmetry display, the answer on the question on existence of Higgs boson, and possibly a new insight on origin of Dark Matter ([1], [2]) but permits one to reconsider the problems of an unique description of variety of elementary particles and all known interactions. If the above hopes are true the development of higher-spin (HS) field theory in view of its close relation to superstring theory on constant curvature spaces, which operates with an infinite set of massive and massless bosonic and fermionic HS fields subject to multi-row Young tableaux (YT) $Y(s_1, \dots, s_k)$, $k \geq 1$ (see for a review, [3]–[7]) seems by actual one. Corresponding description of such theories, having as the final aim the Lagrangian form, requires quite modern and complicated group-theoretic tools which are connected with a construction of different representations of algebras and superalgebras underlying above theories. Whereas for Lie (super)algebra case relevant for HS fields on flat spaces the finding and structure of mentioned objects like Verma modules and generalized Verma modules [8], [9] are rather understandable, an analogous situation with non-linear algebraic and superalgebraic structures which corresponds to HS fields on AdS spaces has not been classified to present with except for the case of totally-symmetric bosonic [10]–[12] and fermionic [13], [14] HS fields. The paper propose the results of regular method for Verma module constructing and Fock space realization for quadratic algebras, whose negative (equivalently positive) root

vectors in Cartan-like triangular decomposition are entangled due to presence of the special parameter r being by the inverse square of AdS-radius vanishing in the flat space limit. The obtained objects permit to find Lagrangian formulations (LFs) for free integer HS fields on d -dimensional AdS-space with $Y(s_1, \dots, s_k)$ in Fronsdal [15] metric-like formalism within BFV-BRST procedure [16], [17] (known at present as BRST construction) as a starting point for an interacting HS field theory in the framework of conventional Quantum Field Theory. The application of the BRST construction to free HS field theory on AdS spaces consists in 4 steps and presents a solution of the *problem inverse* to that of the method [16] (as in the case of string field theory [18]) and reflects one more side of the BV–BFV duality concept [19]–[21]. First, the conditions that determine the representations with given mass and spin are regarded as a topological (i.e. without Hamiltonian) gauge system of mixed-class operator constraints o_I , $I = 1, 2, \dots$, in an auxiliary Fock space \mathcal{H} . Second, the whole system of o_I which form quadratic commutator algebra should be additively converted (see [22], [23] for the development of conversion methods) in deformed (in power of parameter r) algebra of O_I : $O_I = o_I + o'_I$ determined on wider Fock space, $\mathcal{H} \otimes \mathcal{H}'$ with first-class constraints O_α , $O_\alpha \subset O_I$. Third, the, Hermitian nilpotent BFV-BRST operator Q' for non-linear algebra of converted operators O_I which contains the BFV-BRST operator Q for only subsystem of O_α should be found. Fourth, the Lagrangian \mathcal{L} for given HS field through a scalar product $\langle | \rangle$ in total Hilbert space \mathcal{H}_{tot} like $\mathcal{L} \sim \langle \phi | Q | \phi \rangle$, to be invariant with respect to (reducible) gauge transformations $\delta|\phi\rangle = Q|\Lambda\rangle$ with $|\phi\rangle$ containing initial and auxiliary HS fields is constructed in such a way that the corresponding equations of motion reproduce the initial constraints.

The application of above algorithm for bosonic [24], [25] and fermionic [26]–[28] HS fields on flat spaces did not meet the problems in rather complicated second and third steps due to linear Lie structure of initial constraints o_I algebra, $[o_I, o_J] = f_{IJ}^K o_K$ for structure constants f_{IJ}^K . Indeed, for the same algebra of additional parts o'_I it is sufficient to use the construction of Verma module (VM) for integer HS symmetry algebra $sp(2k)$ [29] which is in one-to-one correspondence with Lorentz $so(1, d-1)$ algebra unitary irreducible representations subject to $Y(s_1, \dots, s_k)$, $k \leq \left[\frac{d}{2}\right]$ due to Howe duality [30]. Then an oscillator realization of the symplectic algebra $sp(2k)$ in Fock space \mathcal{H}' [29] represents a polynomial form as compared to totally-symmetric HS fields on AdS space [11]–[14] where, (super)algebras of o_I are non-linear, not coinciding with ones for o'_I . The problem of the same complication arises in constructing of BFV-BRST operator Q' for (super)algebra of converted constraints O_I which in transiting to AdS-space does not have the form to be quadratic in ghost coordinates \mathcal{C} and requires the full study of algebraic relations starting from Jacobi identities resolution (see [31] for details of finding BRST operator Q' in question and ones for classical quadratic algebras [32]).

To be complete, note the details of Lagrangian description of mixed-symmetry HS tensors on (A)dS backgrounds were studied in "frame-like" formulation in [33]–[36] whereas the LF for the mixed-symmetry bosonic fields with off-shell traceless constraints in the case of (anti)-de-Sitter case are recently known for the Young tableaux with two rows [37], [38]. The aspects of $SO(N)$ spinning particles dynamics after applying of Dirac quantization procedure to particle's first-class constraints system which produces the dynamics of HS fields on constant curvature spaces were studied in [39]. At last, the various aspects of mixed-symmetry HS fields Lagrangian dynamics on Minkowski space were discussed in [40], [41] and recently for interacting mixed-symmetry HS fields on AdS-spaces in [42], [43].

Present paper is devoted to the solution of the following problems:

- (i) derivation of HS symmetry algebra for bosonic HS fields in d -dimensional AdS space subject to arbitrary YT $Y(s_1, \dots, s_k)$;
- (ii) development of a method of Verma module construction for a non-linear HS symmetry algebra for YT with two rows $Y(s_1, s_2)$ and to the oscillator realization for given non-

linear algebra as formal power series in creation and annihilation operators of corresponding Heisenberg algebra;

(iii) construction of unconstrained LF for free bosonic HS fields on AdS-space with $Y(s_1, s_2)$.

The paper is organized as follows. In Section 2, we examine the bosonic HS fields that includes, first, the derivation of HS symmetry algebra $\mathcal{A}(Y(k), AdS_d)$ for HS fields subject to Young tableaux with arbitrary number of rows k in Subsection 2.1, second, an auxiliary proposition that permits one to find the forms of deformation of general commutator polynomial algebras under additive conversion procedure in Subsection 2.2. In Section 3 we derive explicitly HS symmetry algebra of additional parts for HS fields with 2 families of indices, formulate and solve the problem of Verma module construction for algebra $\mathcal{A}'(Y(2), AdS_d)$ of additional parts \mathcal{O}'_I in Subsection 3.1. We find Fock space realization for the algebra $\mathcal{A}'(Y(2), AdS_d)$ in Subsection 3.2. In Section 4, we derive an explicit form for non-linear algebra $\mathcal{A}_c(Y(2), AdS_d)$ of converted operators \mathcal{O}_I , and present for it an expression for BFV-BRST operator in Subsection 4.2. and develop the unconstrained Lagrangian formulation for bosonic HS fields with two-rows Young tableaux $Y(s_1, s_2)$ in Subsection 4.3. In Conclusion, we summarize the results of the work and discuss some open problems. Finally, in Appendix A we prove the proposition on additive conversion for polynomial algebras.

2. HS fields in AdS spaces with integer spin

In the section we derive numbers of special HS symmetry non-linear algebras which encode mixed-symmetry tensor fields as the elements of AdS group irreducible representations with generalized spin $\mathbf{s} = (s_1, \dots, s_k)$ and mass m on AdS_d -space-time. We consider the problem of Verma module construction for one of them and solve it explicitly for non-linear algebra with two-rows Young tableaux. The construction of the Fock space representation for the non-linear algebra with found Verma module finishes the solution of the problem there.

2.1. HS symmetry algebra $\mathcal{A}(Y(k), AdS_d)$ for mixed-symmetry tensor fields with $Y(s_1, \dots, s_k)$

A massive generalized integer spin $\mathbf{s} = (s_1, \dots, s_k)$, $(s_1 \geq s_2 \geq \dots \geq s_k > 0, k \leq [d/2])$, AdS group irreducible representation in an AdS_d space is realized in a space of mixed-symmetry tensors, $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} \equiv \Phi_{\mu_1^1 \dots \mu_{s_1}^1, \mu_1^2 \dots \mu_{s_2}^2, \dots, \mu_1^k \dots \mu_{s_k}^k}(x)$ to be corresponding to a Young tableaux

$$\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} \longleftrightarrow \begin{array}{|c|c|c|c|c|c|c|c|c|} \hline \mu_1^1 & \mu_2^1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \mu_{s_1}^1 \\ \hline \mu_1^2 & \mu_2^2 & \cdot & \cdot & \cdot & \cdot & \cdot & \mu_{s_2}^2 & \\ \hline \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \\ \hline \mu_1^k & \mu_2^k & \cdot & \cdot & \cdot & \cdot & \mu_{s_k}^k & & \\ \hline \end{array}, \quad (1)$$

subject to the Klein-Gordon (2), divergentless (3), traceless (4) and mixed-symmetry equations (5) [for $\beta = (2; 3; \dots; k+1) \iff (s_1 > s_2; s_1 = s_2 > s_3; \dots; s_1 = s_2 = \dots = s_k)$] [44]:

$$[\nabla^2 + r[(s_1 - \beta - 1 + d)(s_1 - \beta) - \sum_{i=1}^k s_i] + m^2] \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = 0, \quad (2)$$

$$\nabla^{\mu_{l_i}^i} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = 0, \quad i, j = 1, \dots, k; l_i, m_i = 1, \dots, s_i, \quad (3)$$

$$g^{\mu_{l_i}^i, \mu_{m_i}^i} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = g^{\mu_{l_i}^i, \mu_{m_j}^j} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}} = 0, \quad l_i < m_i, \quad (4)$$

$$\Phi_{(\mu^1)_{s_1}, \dots, \{(\mu^i)_{s_i}, \underbrace{\mu_1^j \dots \mu_{l_j}^j}_{\text{do not include in symmetrization}}, \dots, \mu_{s_j}^j, \dots, (\mu^k)_{s_k}\}} = 0, \quad i < j, 1 \leq l_j \leq s_j, \quad (5)$$

where the brackets below denote that the indices inside it do not include in symmetrization, i.e. the symmetrization concerns only indices $(\mu^i)_{s_i}, \mu_{l_j}^j$ in $\{(\mu^i)_{s_i}, \underbrace{\mu_1^j \dots \mu_{l_j}^j}_{\text{do not include in symmetrization}}, \dots, \mu_{s_j}^j\}$.

To obtain HS symmetry algebra (of o_I) for a description of all integer spin HS fields, we in a standard manner introduce a Fock space \mathcal{H} , generated by k pairs of bosonic creation $a_{\mu^i}^i(x)$ and annihilation $a_{\nu^j}^{j+}(x)$ operators, $i, j = 1, \dots, k, \mu^i, \nu^j = 0, 1, \dots, d-1$

$$[a_{\mu^i}^i, a_{\nu^j}^{j+}] = -g_{\mu^i \nu^j} \delta^{ij}, \quad \delta^{ij} = \text{diag}(1, 1, \dots, 1), \quad (6)$$

and a set of constraints for an arbitrary string-like vector $|\Phi\rangle \in \mathcal{H}$ which we call as the basic vector,

$$|\Phi\rangle = \sum_{s_1=0}^{\infty} \sum_{s_2=0}^{s_1} \cdots \sum_{s_k=0}^{s_{k-1}} \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}}(x) \prod_{i=1}^k \prod_{l_i=1}^{s_i} a_i^{+\mu_{l_i}^i} |0\rangle, \quad (7)$$

$$\tilde{l}_0 |\Phi\rangle = (l_0 + \tilde{m}_b^2 + r((g_0^1 - 2\beta - 2)g_0^1 - \sum_{i=2}^k g_0^i)) |\Phi\rangle = 0, \quad l_0 = [D^2 - r \frac{d(d-2(k+1))}{4}], \quad (8)$$

$$(l^i, l^{ij}, t^{i_1 j_1}) |\Phi\rangle = (-i a_{\mu}^i D^{\mu}, \frac{1}{2} a_{\mu}^i a^{j\mu}, a_{\mu}^{i_1+} a^{j_1 \mu}) |\Phi\rangle = 0, \quad i \leq j; i_1 < j_1, \quad (9)$$

with number particles operators, central charge, covariant derivative in \mathcal{H} respectively,

$$g_0^i = -\frac{1}{2} \{a_{\mu}^{i+}, a^{\mu i}\}, \quad \tilde{m}_b^2 = m^2 + r\beta(\beta + 1), \quad (10)$$

$$D_{\mu} = \partial_{\mu} - \omega_{\mu}^{ab}(x) \left(\sum_i a_{ia}^{+} a_{ib} \right), \quad a_i^{\mu(+)}(x) = e_a^{\mu}(x) a_i^{a(+)}(x), \quad (11)$$

where e_a^{μ} , ω_{μ}^{ab} are vielbein and spin connection for tangent indices $a, b = 0, 1, \dots, d-1$. Operator D_{μ} is equivalent in its action in \mathcal{H} to the covariant derivative ∇_{μ} [with d'Alembertian $D^2 = (D_a + \omega_{ba}^b) D^a$]. The set of $k(k+1)$ primary constraints (8), (9) with $\{o_{\alpha}\} = \{\tilde{l}_0, l^i, l^{ij}, t^{i_1 j_1}\}$ are equivalent to Eqs. (2)–(5) for all admissible values of spins and for the field $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}}$ with fixed spin $\mathbf{s} = (s_1, s_2, \dots, s_k)$ if in addition to Eqs. (8), (9) we add k constraints with g_0^i ,

$$g_0^i |\Phi\rangle = (s_i + \frac{d}{2}) |\Phi\rangle. \quad (12)$$

The requirement of closedness of the algebra with o_{α} with respect to $[\ , \]$ -multiplication leads to enlargement of o_{α} by adding the operators g_0^i and hermitian conjugated operators o_{α}^{+} ,

$$(l^{i+}, l^{ij+}, t^{i_1 j_1+}) = (-i a_{\mu}^{i+} D^{\mu}, \frac{1}{2} a_{\mu}^{i+} a^{j\mu+}, a_{\mu}^{i_1+} a^{j_1 \mu+}), \quad i \leq j; i_1 < j_1, \quad (13)$$

with respect to scalar product on \mathcal{H} ,

$$\begin{aligned} \langle \Psi | \Phi \rangle &= \int d^d x \sqrt{|g|} \sum_{i=1}^k \sum_{s_i=0}^{s_{i-1}} \sum_{j=1}^k \sum_{p_j=0}^{s_{j-1}} \langle 0 | \prod_{j=1}^k \prod_{m_j=1}^{p_j} a_j^{\nu_{m_j}^j} \Psi_{(\nu^1)_{p_1}, (\nu^2)_{p_2}, \dots, (\nu^k)_{p_k}}^* (x) \\ &\quad \times \Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}}(x) \prod_{i=1}^k \prod_{l_i=1}^{s_i} a_i^{+\mu_{l_i}^i} |0\rangle, \quad \text{for } s_{-1}, p_{-1} = \infty. \end{aligned} \quad (14)$$

¹ such choice of the oscillators corresponds to the case of symmetric basis, whereas there exists another realization of auxiliary Fock space generated by the fermionic oscillators (antisymmetric basis) $\hat{a}_{\mu^m}^m(x)$, $\hat{a}_{\nu^n}^{n+}(x)$ with anticommutation relations, $\{\hat{a}_{\mu^m}^m, \hat{a}_{\nu^n}^{n+}\} = -g_{\mu^m \nu^n} \delta^{mn}$, for $m, n = 1, \dots, s_1$, and develop the procedure below following to the lines of Ref. [45] for totally antisymmetric tensors for $s_1 = s_2 = \dots = s_k = 1$.

² operators $a_i^{a_i}$, $a_j^{b_j+}$ satisfy to usual for $R^{1,d-1}$ -space commutation relations $[a_i^{a_i}, a_j^{b_j+}] = -\eta^{a_i b_i} \delta_{ij}$ for $\eta^{ab} = \text{diag}(+, -, \dots, -)$

Table 1. HS symmetry non-linear algebra $\mathcal{A}_m(Y(k), AdS_d)$.

| $[\downarrow, \rightarrow]$ | $t^{i_1 j_1}$ | $t_{i_1 j_1}^+$ | l_0 | l^i | l^{i+} | $l^{i_1 j_1}$ | $l^{i_1 j_1+}$ |
|-----------------------------|-------------------------------------|-------------------------------------|-----------------------|--|---|--|--|
| $t^{i_2 j_2}$ | $A^{i_2 j_2, i_1 j_1}$ | $B^{i_2 j_2, i_1 j_1}$ | 0 | $l^{j_2} \delta^{i_2 i}$ | $-l^{i_2+} \delta^{j_2 i}$ | $l^{\{j_1 j_2\} \delta^{i_1\}}$ | $-l^{i_2 \{i_1+ \delta^{j_1\}} j_2}$ |
| $t_{i_2 j_2}^+$ | $-B^{i_1 j_1, i_2 j_2}$ | $A_{i_1 j_1, i_2 j_2}^+$ | 0 | $l_{i_2} \delta_{j_2}^i$ | $-l_{j_2}^+ \delta_{i_2}^i$ | $l_{i_2} \{j_1 \delta_{j_2}^{i_1}\}$ | $-l_{j_2} \{j_1+ \delta_{i_2}^{i_1}\}$ |
| l_0 | 0 | 0 | 0 | $-r\mathcal{K}_1^{i+}$ | $r\mathcal{K}_1^i$ | 0 | 0 |
| l^j | $-l^{j_1} \delta^{i_1 j}$ | $-l^{i_1} \delta^{j_1 j}$ | $r\mathcal{K}_1^{j+}$ | $W_b^{j_1}$ | $X_b^{j_1}$ | 0 | $-\frac{1}{2} l^{\{i_1+ \delta^{j_1\}} j}$ |
| l^{j+} | $l^{i_1+} \delta^{j_1 j}$ | $l^{j_1+} \delta^{i_1 j}$ | $-r\mathcal{K}_1^j$ | $-X_b^{ij}$ | $-W_b^{j+}$ | $\frac{1}{2} l^{\{i_1 \delta^{j_1\}} j}$ | 0 |
| $l^{i_2 j_2}$ | $-l^{j_1 \{j_2 \delta^{i_2\}} i_1}$ | $-l^{i_1 \{i_2 \delta^{j_2\}} j_1}$ | 0 | 0 | $-\frac{1}{2} l^{\{i_2 \delta^{j_2\}} i}$ | 0 | $L^{i_2 j_2, i_1 j_1}$ |
| $l_{i_2 j_2}^+$ | $l_{i_2}^{i_1+} \delta_{j_2}^{j_1}$ | $l_{j_1}^+ \delta_{i_2}^{i_1}$ | 0 | $\frac{1}{2} l_{i_2}^+ \delta_{j_2}^i$ | 0 | $-L^{i_1 j_1, i_2 j_2}$ | 0 |
| g_0^j | $-F^{i_1 j_1, j}$ | $F^{i_1 j_1, j+}$ | 0 | $-l^i \delta^{ij}$ | $l^{i+} \delta^{ij}$ | $-l^j \{i_1 \delta^{j_1\}} j$ | $l^j \{i_1+ \delta^{j_1\}} j$ |

This fact will guarantee the Hermiticity of corresponding BFV-BRST operator with taken into account of self-conjugated operators, $(l_0^+, g_0^{i+}) = (l_0, g_0^i)$ (therefore the reality of Lagrangian \mathcal{L}) for the system of all operators $\{o_\alpha, o_\alpha^+, g_0^i\}$. We call the algebra of these operators the *integer higher-spin symmetry algebra in AdS space with a Young tableaux having k rows*³ and denote it as $\mathcal{A}(Y(k), AdS_d)$.

The maximal Lie subalgebra of operators $l^{ij}, t^{i_1 j_1}, g_0^i, l^{ij+}, t^{i_1 j_1+}$ is isomorphic to symplectic algebra $sp(2k)$ (see, [29] for details and we will refer on it later as $sp(2k)$) whereas the only nontrivial quadratic commutators in $\mathcal{A}(Y(k), AdS_d)$ are due to operators with D_μ : l^i, \tilde{l}_0, l^{i+} . For the aim of LF construction it is enough to have a simpler, without central charge \tilde{m}_b^2 (so called *modified* HS symmetry algebra $\mathcal{A}_m(Y(k), AdS_d)$), with operator l_0 (8) instead of \tilde{l}_0 , so that AdS-mass term, $\tilde{m}_b^2 + r((g_0^1 - 2\beta - 2)g_0^1 - \sum_{i=2}^k g_0^i)$, will be restored as usual later within conversion procedure and properly construction of LF.

Algebra $\mathcal{A}_m(Y(k), AdS_d)$ of the operators o_I from the Hamiltonian analysis of the dynamical systems viewpoint contains 1 first-class constraint l_0 , $2k$ differential l_i, l_i^+ and $2k^2$ algebraic $t^{i_1 j_1}, t_{i_1 j_1}^+, l^{ij}, l_{ij}^+$ second-class constraints o_a and operators g_0^i , composing an invertible matrix $\Delta_{ab}(g_0^i)$ for topological gauge system because of

$$[o_a, o_b] = f_{ab}^c o_c + f_{ab}^{cd} o_c o_d + \Delta_{ab}(g_0^i), [o_a, l_o] = f_{a[l_0]}^c o_c + f_{a[l_0]}^{cd} o_c o_d, . \quad (15)$$

Here $f_{ab}^c, f_{ab}^{cd}, f_{a[l_0]}^c, f_{a[l_0]}^{cd}, \Delta_{ab}$ are antisymmetric with respect to permutations of lower indices constant quantities and the operators $\Delta_{ab}(g_0^i)$ form the non-vanishing $2k(k+1) \times 2k(k+1)$ matrix $\|\Delta_{ab}\|$ in the Fock space \mathcal{H} on the surface $\Sigma \subset \mathcal{H}$: $\|\Delta_{ab}\|_\Sigma \neq 0$, which is determined by the equations, $o_\alpha|\Phi\rangle = 0$. The set of o_I satisfies the non-linear relations (additional to ones for $sp(2k)$) given by the multiplication table 1.

First note that, in the table 1 we did not include the columns with $[\downarrow, \rightarrow]$ -products of all o_I with g_0^j which may be obtained from the rows with g_0^j as follows: $[b, o_I]^+ = -[o_I^+, b]$, with account of closedness of HS algebra with respect to Hermitian conjugation. Second the operators $t^{i_2 j_2}, t_{i_2 j_2}^+$ satisfy by the definition the properties

$$(t^{i_2 j_2}, t_{i_2 j_2}^+) \equiv (t^{i_2 j_2}, t_{i_2 j_2}^+) \theta^{j_2 i_2}, \theta^{j_2 i_2} = (1, 0) \text{ for } (j_2 > i_2, j_2 \leq i_2) \quad (16)$$

³ one should not confuse the term "higher-spin symmetry algebra" using here for free HS formulation with the algebraic structure known as "higher-spin algebra" (see, e.g. Ref.[46]) arising to describe the HS interactions

with Heaviside θ -symbol⁴ θ^{ji} . Third, the products $B_{i_1 j_1}^{i_2 j_2}, A^{i_2 j_2, i_1 j_1}, F^{i_1 j_1, i}, L^{i_2 j_2, i_1 j_1}$ are determined by the explicit relations,

$$B^{i_2 j_2}_{i_1 j_1} = (g_0^{i_2} - g_0^{j_2})\delta_{i_1}^{i_2}\delta_{j_1}^{j_2} + (t_{j_1}^{j_2}\theta^{j_2}_{j_1} + t_{j_1}^{j_2+}\theta_{j_1}^{j_2})\delta_{i_1}^{i_2} - (t_{i_1}^{+i_2}\theta^{i_2}_{i_1} + t_{i_1}^{i_2}\theta_{i_1}^{i_2})\delta_{j_1}^{j_2}, \quad (17)$$

$$A^{i_2 j_2, i_1 j_1} = t^{i_1 j_2}\delta^{i_2}_{j_1} - t^{i_2 j_1}\delta^{i_1}_{j_2}, \quad F^{i_2 j_2, i} = t^{i_2 j_2}(\delta^{j_2 i} - \delta^{i_2 i}), \quad (18)$$

$$L^{i_2 j_2, i_1 j_1} = \frac{1}{4}\left\{\delta^{i_2 i_1}\delta^{j_2 j_1}\left[2g_0^{i_2}\delta^{i_2 j_2} + g_0^{i_2} + g_0^{j_2}\right] - \delta^{j_2}\{i_1\left[t_{j_1}^{j_2}\theta^{i_2}_{j_1}\right] + t^{i_2 j_1}\} + \theta_{j_1}^{i_2}\right\} \\ - \delta^{i_2}\{i_1\left[t_{j_1}^{j_2}\theta^{j_2}_{j_1}\right] + t^{j_2 j_1}\} + \theta_{j_1}^{j_2}\}. \quad (19)$$

They obeys the obvious additional properties of antisymmetry and Hermitian conjugation,

$$A^{i_2 j_2, i_1 j_1} = -A^{i_1 j_1, i_2 j_2}, \quad A_{i_1 j_1, i_2 j_2}^+ = (A_{i_1 j_1, i_2 j_2})^+ = t_{i_2 j_1}^+ \delta_{j_2 i_1} - t_{i_1 j_2}^+ \delta_{i_2 j_1}, \quad (20)$$

$$(L^{i_2 j_2, i_1 j_1})^+ = L^{i_1 j_1, i_2 j_2}, \quad F^{i_2 j_2, i+} = (F^{i_2 j_2, i})^+ = t^{i_2 j_2+}(\delta^{j_2 i} - \delta^{i_2 i}), \quad (21)$$

$$B^{i_2 j_2}_{i_1 j_1}^+ = (g_0^{i_2} - g_0^{j_2})\delta_{i_1}^{i_2}\delta_{j_1}^{j_2} + (t_{j_1}^{j_2}\theta^{j_2}_{j_1} + t_{j_1}^{j_2+}\theta_{j_1}^{j_2})\delta_{i_1}^{i_2} - (t_{i_1}^{i_2}\theta^{i_2}_{i_1} + t_{i_1}^{i_2+}\theta_{i_1}^{i_2})\delta_{j_1}^{j_2}. \quad (22)$$

Fourth, the independent quantities $\mathcal{K}_1^k, W_b^{ki}, X_b^{ki}$ in the table 1 are quadratic in O_I ,

$$W_b^{ij} = [l^i, l^j] = 2r\left[(g_0^j - g_0^i)l^{ij} - \sum_m (t^{m[j}\theta^{j]m} + t^{[jm+}\theta^{m]j})l^i\right], \quad (23)$$

$$\mathcal{K}_1^k = r^{-1}[l_0, l^{k+}] = \left(4\sum_i l^{ki+}l^i + l^{k+}(2g_0^k - 1) - 2(\sum_i l^{i+}t^{+ik}\theta^{ki} + l^{i+}t^{ki}\theta^{ik})\right), \quad (24)$$

$$X_b^{ij} = \{l_0 + r(K_0^{0i} + \sum_{l=i+1}^k \mathcal{K}_0^{il} + \sum_{l=1}^{i-1} \mathcal{K}_0^{li})\}\delta^{ij} \\ - r[4\sum_l l^{jl+}l^i - \sum_{l=1}^{j-1} t^{lj+}t^{li} - \sum_{l=i+1}^k t^{il+}t^{jl} - \sum_{l=j+1}^{i-1} t^{li}t^{jl} + (g_0^j + g_0^i - j - 1)t_{ji}]\theta^{ij} \\ - r[4\sum_l l^{jl+}l^i - \sum_{l=1}^{i-1} t^{lj+}t^{li} - \sum_{l=j+1}^k t^{il+}t^{jl} - \sum_{l=i+1}^{j-1} t^{il+}t^{lj+} + t_{ij}^+(g_0^j + g_0^i - i - 1)]\theta^{ji}. \quad (25)$$

In writing (25) we have used the quantities $K_0^{0i}, i, j = 1, \dots, k$, \mathcal{K}_0^{ij} composing a Casimir operator $\mathcal{K}_0(k)$ for $sp(2k)$ algebra

$$\mathcal{K}_0(k) = \sum_i K_0^{0i} + 2\sum_{i,j} \mathcal{K}_0^{ij}\theta^{ji} = \sum_i ((g_0^i)^2 - 2g_0^i - 4l^{ii+}l^{ii}) + 2(\sum_{i=1}^k \sum_{j=i+1}^k (t_{ij}^+ t^{ij} - 4l_{ij}^+ l^{ij} - g_0^j)). \quad (26)$$

Algebra $\mathcal{A}_m(Y(k), AdS_d)$ maybe considered as non-linear deformation in power of r of the integer HS symmetry algebra in Minkowski space $\mathcal{A}(Y(k), R^{1,d-1})$ [29] as follows,

$$\mathcal{A}_m(Y(k), AdS_d) = \mathcal{A}(Y(k), R^{1,d-1})(r) = (T^k \oplus T^{k*} \oplus l_0)(r) \oplus sp(2k), \quad (27)$$

for k -dimensional commutative (on $R^{1,d-1}$) algebra $T^k = \{l_i\}$, its dual $T^{k*} = \{l^{i+}\}$ and represents the semidirect sum of the symplectic algebra $sp(2k)$ [as an algebra of internal derivations of $(T^k \oplus T^{k*})$]. For HS fields with single spin s_1 (for $k = 1$) and with two-component spin (s_1, s_2) (for $k = 2$) the algebra $\mathcal{A}_m(Y(k), AdS_d)$ coincides respectively with known HS symmetry algebras on AdS spaces given in [12] and in [31].

Now, we should to use the results of an special proposition in order to proceed to the conversion procedure of the algebra of O_I to get the algebra of O_I with only first-class constraints.

⁴ there are no summation with respect to the indices i_2, j_2 in the Eqs.(16), the figure brackets for the indices i_1, i_2 in the quantity $A^{\{i_1 B^{i_2\}i_3\}\theta^{i_3 i_2}}$ mean the symmetrization $A^{\{i_1 B^{i_2\}i_3\}\theta^{i_3 i_2}} = A^{i_1 B^{i_2 i_3}\theta^{i_3 i_2}} + A^{i_2 B^{i_1 i_3}\theta^{i_3 i_1}}$ as well as these indices are raising and lowering by means of Euclidian metric tensors $\delta^{ij}, \delta_{ij}, \delta_j^i$

2.2. On additive conversion for polynomial algebras

In this subsection, to solve the problem of additive conversion of non-linear algebras with a subset of 2nd class constraints, we need to use some important statement which based on the following (see Ref.[14] for detailed description)

Definition: A non-linear commutator algebra \mathcal{A} of basis elements o_I , $I \in \Delta$ (with Δ to be finite or infinite set of indices) is called a **polynomial algebra of order n** , $n \in \mathbf{N}$, if a set of $\{o_I\}$ is subject to n -th order polynomial commutator relations:

$$[o_I, o_J] = F_{IJ}^K(o) o_K, \quad F_{IJ}^K = f_{IJ}^{(1)K} + \sum_{n=2}^{\infty} f_{IJ}^{(n)K_1 \dots K_{n-1}K} \prod_{i=1}^{n-1} o_{K_i},$$

$$f_{IJ}^{(n)K_1 \dots K_n} \neq 0, \quad \text{and} \quad f_{IJ}^{(k)K_1 \dots K_n K_{n+1} \dots K_k} = 0, \quad k > n, \quad (28)$$

with structural coefficients $f_{IJ}^{(n)K_1 \dots K_{n-1}K}$, to be antisymmetric with respect to permutations of lower indices, $f_{IJ}^{(n)K_1 \dots K_n} = -f_{JI}^{(n)K_1 \dots K_n}$.

Now, we may to formulate the basic statement in the subsection 2.2 in the form of

Proposition: Let \mathcal{A} is the polynomial algebra of order n of basis elements o_I determined in Hilbert space \mathcal{H} . Then, for a set \mathcal{A}' of elements o'_I given in a new Hilbert space \mathcal{H}' ($\mathcal{H} \cap \mathcal{H}' = \emptyset$) and commuting with o_I , and for a direct sum of sets $\mathcal{A}_c = \mathcal{A} + \mathcal{A}'$ of the operators, $O_I = o_I + o'_I$ given in the tensor product $\mathcal{H} \otimes \mathcal{H}'$ from the requirement to be in involution relations,

$$[O_I, O_J] = F_{IJ}^K(o', O) O_K, \quad (29)$$

it follows the sets of $\{o'_I\}$, $\{O_I\}$ form respectively the polynomial commutator algebra \mathcal{A}' of order n and the non-linear commutator algebra \mathcal{A}_c with composition laws:

$$[o'_I, o'_J] = f_{IJ}^{(1)K_1} o'_{K_1} + \sum_{m=2}^n (-1)^{m-1} f_{IJ}^{(m)K_1 \dots K_1} \prod_{s=1}^m o'_{K_s}, \quad (30)$$

$$[O_I, O_J] = \left(f_{IJ}^{(1)K} + \sum_{m=2}^n F_{IJ}^{(m)K}(o', O) \right) O_K. \quad (31)$$

The structural functions $F_{IJ}^{(m)k}(o', O)$ in (30) are constructed with respect to known from the Eqs. (28) coefficients $f_{IJ}^{(m)K_1 \dots K_m} \equiv f_{IJ}^{K_1 \dots K_m}$ as follows

$$F_{IJ}^{(m)K} = f_{IJ}^{K_1 \dots K_m} \prod_{p=1}^{m-1} O_{K_p} + \sum_{s=1}^{m-1} (-1)^s f_{ij}^{K_s \dots K_1 K_{s+1} \dots K_m} \prod_{p=1}^s o'_{K_p} \prod_{l=s+1}^{m-1} O_{K_l}, \quad \text{where}$$

$$f_{ij}^{K_s \dots K_1 K_{s+1} \dots K_m} = f_{ij}^{K_s \dots K_1 K_{s+1} \dots K_m} + f_{ij}^{K_s \dots K_{s+1} K_1 K_{s+2} \dots K_m} + \dots +$$

$$f_{ij}^{K_{s+1} K_s \dots K_1 K_{s+2} \dots K_m} + \left(f_{ij}^{K_{s+1} K_s \dots K_{s+2} K_1 K_{s+3} \dots K_m} + \dots + f_{ij}^{K_{s+1} K_{s+2} K_s \dots K_1 K_{s+3} \dots K_l} \right) + \dots + f_{ij}^{K_{s+1} \dots K_m K_s \dots K_1}, \quad (32)$$

where the sum in the Eq.(32) contains $\frac{m!}{s!(m-s)!}$ terms with all the possible ways of the arrangement the indices (K_{s+1}, \dots, K_m) among the indices (K_s, \dots, K_1) in $f_{ij}^{K_s \dots K_1 K_{s+1} \dots K_m}$ without changing the separate orderings of the indices K_{s+1}, \dots, K_m and K_s, \dots, K_1 ⁵.

⁵ We do not consider here the case of polynomial superalgebra for which the proposition may be easily generalized with introducing corresponding sign factors in the Eqs. (30)–(32) with the same number of summands to use it for fermionic HS fields on AdS space

The validity of the proposition is verified in the Appendix A. Turning to the structure of algebra \mathcal{A}_c note that in contrary to \mathcal{A} and \mathcal{A}' we call it as *non-homogeneous polynomial algebra of order n* due to form of relations (32) (see footnote 13 in the Appendix A for the comments). For Lie algebra case ($n = 1$) the structures of the algebras \mathcal{A} , \mathcal{A}' and \mathcal{A}_c coincides as it then used, e.g. for the integer spin HS symmetry algebra $\mathcal{A}(Y(k), R^{1,d-1})$ [29]. For quadratic algebras ($n=2$) the algebraic relations for \mathcal{A} , \mathcal{A}' and \mathcal{A}_c do not coincide with each other due to structural functions $f_{IJ}^{(2)K_1K_2}$ presence that was firstly shown for the algebra $\mathcal{A}(Y(1), AdS_d)$ for totally-symmetric HS tensors on AdS space in [12] and having the form,

$$[o_I, o_J] = f_{IJ}^{(1)K_1} o_{K_1} + f_{IJ}^{(2)K_1K_2} o_{K_1} o_{K_2}, \quad [o'_I, o'_J] = f_{IJ}^{(1)K_1} o'_{K_1} - f_{IJ}^{(2)K_2K_1} o'_{K_1} o'_{K_2}, \quad (33)$$

$$[O_I, O_J] = (f_{IJ}^{(1)K} + F_{IJ}^{(2)K}(o', O)) O_K, \quad F_{IJ}^{(2)K} = f_{IJ}^{(2)K_1K} O_{K_1} - (f_{IJ}^{(2)K_1K} + f_{IJ}^{(2)KK_1}) o'_{K_1}. \quad (34)$$

Relations (33), (34) are sufficient to determine the form of the multiplication laws for the additional parts o'_I algebra $\mathcal{A}'(Y(k), AdS_d)$ and for converted operators O_I algebra $\mathcal{A}_c(Y(k), AdS_d)$.

As the new result we write down the explicit form for the cubic commutator algebras \mathcal{A}' , \mathcal{A}_c

$$\begin{aligned} [o'_I, o'_J] &= f_{IJ}^{(1)K_1} o'_{K_1} - f_{IJ}^{(2)K_2K_1} o'_{K_1} o'_{K_2} + f_{IJ}^{(3)K_3K_2K_1} o'_{K_1} o'_{K_2} o'_{K_3}, \\ [O_I, O_J] &= \left(f_{IJ}^{(1)K} + \sum_{l=2}^n F_{IJ}^{(l)K}(o', O) \right) O_K, \end{aligned} \quad (35)$$

$$\begin{aligned} F_{IJ}^{(3)K} &= f_{IJ}^{(3)K_1K_2K} O_{K_1} O_{K_2} - (f_{IJ}^{(3)K_1K_2K} + f_{IJ}^{(3)K_2K_1K}) o'_{K_1} O_{K_2} \\ &\quad + (f_{IJ}^{(3)K_2K_1K} + f_{IJ}^{(3)K_2KK_1} + f_{IJ}^{(3)KK_2K_1}) o'_{K_1} o'_{K_2}, \end{aligned} \quad (36)$$

if the commutator relations for the initial algebra \mathcal{A} given by the Eqs.(28) for $n = 3$.

3. Auxiliary HS symmetry algebra $\mathcal{A}'(Y(2), AdS_d)$

The procedure of additive conversion for non-linear HS symmetry algebra $\mathcal{A}(Y(k), AdS_d)$ of the operators o_I implies finding, first, the explicit form of the algebra $\mathcal{A}'(Y(k), AdS_d)$ of the additional parts o'_I , second, the representation of $\mathcal{A}'(Y(k), AdS_d)$ in terms of some appropriate Heisenberg algebra elements acting in a new Fock space \mathcal{H}' . Structure of the non-linear commutators of the initial algebra leads to necessity to convert all the operators o_I to construct unconstrained LF for given HS field $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}, \dots, (\mu^k)_{s_k}}$.

Considering here the case of $k = 2$ family of indices only in the initial HS field $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}}$ (the general case of algebra $\mathcal{A}'(Y(k), AdS_d)$ is discussed in [47]) we see the former step is based on a determination of a multiplication table $\mathcal{A}'(Y(2), AdS_d)$ of operators o'_I following to the form of the algebra $\mathcal{A}(Y(k), AdS_d)$ for $k = 2$ given by the table 1 and Eqs.(33).

As the result, the searched composition law for $\mathcal{A}'(Y(2), AdS_d)$ is the same as for the algebra $\mathcal{A}(Y(2), AdS_d)$ in its linear Lie part, i.e. for $sp(4)$ subalgebra of elements $(l'^{ij}, l'^{ij+}, t'^{12}, t'^{12+}, g_0'^i)^6$, and is differed in the non-linear part of the Table 1, determined by the isometry group elements l'_i, l'_j, l'_0 . The corresponding non-linear submatrix of the multiplication matrix for $\mathcal{A}'(Y(2), AdS_d)$ has the form given by the Table 2, Here the functions $\mathcal{K}_1'^{i+}, \mathcal{K}_1'^i, W_b'^{ji}, W_b'^{jji+}$ ($X_b'^{ij} - l'_0$) have the same definition as the ones (23)–(25) for initial operators o_I but with opposite sign for $(X_b'^{ij} - l'_0)$ and for $k = 2$:

$$W_b'^{ij} = 2r\epsilon^{ij} \left[(g_0'^2 - g_0'^1) l'^{12} - t'^{12} l'^{11} + t'^{12+} l'^{22} \right], \quad (37)$$

⁶ To turn from general algebra $\mathcal{A}(Y(k), AdS_d)$ to $\mathcal{A}(Y(2), AdS_d)$, we put $\theta^{ij} = \delta^{i2} \delta^{j1}$ and therefore only surviving operators among mixed-symmetry ones are $t'^{ij} = t'^{12}, t'_{ij}^+ = t'_{12}^+$.

Table 2. The non-linear part of algebra $\mathcal{A}'(Y(2), AdS_d)$.

| $[\downarrow, \rightarrow]$ | l'_0 | l'^i | l'^{i+} |
|-----------------------------|-------------------------|-------------------------|----------------------|
| l'_0 | 0 | $r\mathcal{K}_1'^{bi+}$ | $-r\mathcal{K}_1'^i$ |
| l'^j | $-r\mathcal{K}_1'^{j+}$ | $-W_b'^{ij}$ | $X_b'^{ji}$ |
| l'^{j+} | $r\mathcal{K}_1'^j$ | $-X_b'^{ij}$ | $W_b'^{ji+}$ |

$$\mathcal{K}_1'^j = \left(4 \sum_i l'^{ji+} l'^i + l'^{j+} (2g_0'^j - 1) - 2l'^{2+} t'^{+12} \delta^{j1} - 2l'^{1+} t'^{12} \delta^{j2} \right), \quad (38)$$

$$\begin{aligned} X_b'^{ij} &= \{l'_0 - r(K_0'^{0i} + \mathcal{K}_0'^{12})\} \delta^{ij} + r \{ 4 \sum_l l'^{1l+} l'^{l2} + (g_0'^2 + g_0'^1 - 2) t'_{12} \} \delta^{i2} \delta^{j1} \\ &\quad + r \{ 4 \sum_l l'^{l2+} l'^{1l} + t'_{12}^+ (g_0'^2 + g_0'^1 - 2) \} \delta^{i1} \delta^{j2}, \end{aligned} \quad (39)$$

with totally antisymmetric $sp(2)$ -invariant tensor ϵ^{ij} , $\epsilon^{12} = 1$ and operator $\mathcal{K}_0'^{12} = (t_{12}^+ t^{12} - 4l_{12}^+ l^{12} - g_0'^2)$ derived from Casimir operator for $sp(4)$ algebra in the Eqs.(26) for $k = 2$. In turn, the Lie part of the Tables 1 for $k = 2$ is the same as one for bosonic Lie subalgebra in [28] for the following expressions of only non-vanishing operators B'^{12}_{12} , A'^{1212} , $F'^{12,j}$, $F'^{12,j+}$, $L'^{i_1 j_1, i_2 j_2}$

$$B'^{12}_{12} = (g_0'^1 - g_0'^2), \quad A'^{12,12} = 0, \quad F'^{12,j} \equiv F'^j = t'_{12} (\delta^{j2} - \delta^{j1}), \quad (40)$$

$$\begin{aligned} L'^{i_1 j_1, i_2 j_2} &= \frac{1}{4} \left\{ \delta^{i_2 i_1} \delta^{j_2 j_1} \left[2g_0'^{i_2} \delta^{i_2 j_2} + g_0'^{i_2} + g_0'^{j_2} \right] - \delta^{j_2 \{ i_1 \} \left[t'_{12} \delta^{j_1 \} 1 \delta^{i_2 2} + t'_{12}^+ \delta^{j_1 \} 2 \delta^{i_2 1} \right]} \right. \\ &\quad \left. - \delta^{i_2 \{ i_1 \} \left[t'_{12} \delta^{j_2 2} \delta^{j_1 \} 1 + t'_{12}^+ \delta^{j_1 \} 2 \delta^{j_2 1} \right]} \right\}, \end{aligned} \quad (41)$$

being identical to the same operators o_I from the initial algebra $\mathcal{A}(Y(2), AdS_d)$.

Now, we may to sketch the points to find oscillator representation for the elements o'_I of auxiliary HS symmetry algebra.

3.1. Verma module for the quadratic algebra $\mathcal{A}'(Y(2k), AdS_d)$

Here, we following to assumption that generalization of Poincare–Birkhoff–Witt theorem for the second order algebra $\mathcal{A}'(Y(2), AdS_d)$ is true (see on PBW theorem generalization for quadratic algebra [48]), we start to construct Verma module, based on Cartan-like decomposition enlarged from one for $sp(4)$ ($i \leq j$)

$$\mathcal{A}'(Y(2), AdS_d) = \{l'_{ij}^+, t'_{12}^+, l_i'^+\} \oplus \{g_0'^i, l'_0\} \oplus \{l'_{ij}, t'_{12}, l'_i\} \equiv \mathcal{E}_2^- \oplus H_2 \oplus \mathcal{E}_2^+.^7 \quad (42)$$

Note, that in contrast to the case of Lie algebra the element l'_0 does not diagonalize the elements of upper \mathcal{E}_2^- (lower \mathcal{E}_2^+) triangular subalgebra due to quadratic relations (38) (as well as it was for totally-symmetric HS fields on AdS space [10, 12]) and in addition the negative root vectors $l_i'^+, l_j'^+$ do not commute.

Because the Verma module over a semi-simple finite-dimensional Lie algebra g (an induced module $\mathcal{U}(g) \otimes_{\mathcal{U}(b)} |0\rangle_V$ with a vacuum vector $|0\rangle_V$ ⁸) is isomorphic due to PBW theorem as a vector space to a polynomial algebra $\mathcal{U}(g^-) \otimes_C |0\rangle_V$, it is clear that g can be realized by first-order inhomogeneous differential operators acting on these polynomials.

⁷ we may consider $sp(4)$ and generally $sp(2k)$ in Cartan–Weyl basis for unified description, however without loss of generality the basis elements and structure constants of the algebra under consideration will be chosen as in the table 1

⁸ here the signs $\mathcal{U}(g)$, $\mathcal{U}(b)$, $\mathcal{U}(g^-)$ denote the universal enveloping algebras respectively for g , for its Borel subalgebra and for lower triangular subalgebra g^- like \mathcal{E}_2^- in (42)

We consider generalization of *Verma module notion for the case of quadratic algebras* $g(r)$ which present the $g(r)$ -deformation of Lie algebra g in such form that $g(r=0) = g$. Thus, we consider Verma module for such kinds of non-linear algebras, supposing that PBW theorem is valid for $g(r)$ as well that will be proved later by the explicit construction of the Verma module.

Doing so, we consider the quadratic algebra $\mathcal{A}'(Y(2), AdS_d)$, as r -deformation of Lie algebra $\mathcal{A}'(Y(2), R^{1,d-1})$ with use of (27) for $k=2$ (see for details [29]),

$$\mathcal{A}'(Y(2), AdS_d) = (T'^2 \oplus T'^{2*} \oplus l'_0)(r) \oplus sp(4), \quad T'^2 = \{l'_i\}, \quad T'^{2*} = \{l'^{+}_i\}. \quad (43)$$

Corresponding basis vector of Verma module $|\vec{N}(2)\rangle_V$ has, therefore the form according to (42)

$$|\vec{N}(2)\rangle_V = |\vec{n}_{ij}, n_1, p_{12}, n_2\rangle_V \equiv \prod_{i \leq j}^2 (l'^{+}_{ij})^{n_{ij}} \left(\frac{l'_1}{m_1}\right)^{n_1} (t'^{+}_{12})^{p_{12}} \left(\frac{l'_2}{m_2}\right)^{n_2} |0\rangle_V, \quad \mathcal{E}_2^+ |0\rangle_V = 0, \quad (44)$$

with (vacuum) highest weight vector $|0\rangle_V$, non-negative integers n_{ij}, n_l, p_{lm} , and arbitrary constants m_l with dimension of mass.

Here, in contrast to the case of Lie algebra [28] and totally-symmetric HS fields on AdS space [10], [13], the negative root vectors $l'^{+}_1, t'^{+}_{12}, l'^{+}_2$ do not commute, making the vector $|\vec{N}(2)\rangle_V$ by not proper one for the operators t'^{+}_{12}, l'^{+}_2 presenting therefore the essential peculiarities to construct Verma module for $\mathcal{A}'(Y(2), AdS_d)$.

First of all, by the definition the highest weight vector $|0\rangle_V$ is proper for the vectors from the Cartan-like subalgebra H_2 ,

$$(g'_0, l'_0)|0\rangle_V = (h^i, m_0^2)|0\rangle_V, \quad (45)$$

with some real numbers h^1, h^2, m_0 whose values will be determined later in the end of LF construction in. order to Lagrangian equations of motion reproduce the initial AdS group irreps conditions (2)–(5).

Then, we determine the action of the negative root vectors from the subspace \mathcal{E}_2^- on the basis vector $|\vec{N}(2)\rangle_V$ which now do not present explicitly the action of raising operators and reads,

$$l'_{lm}{}^+ |\vec{N}(2)\rangle_V = |\vec{n}_{ij} + \delta_{ij,lm}, \vec{n}_s\rangle_V, \quad (46)$$

$$l'_l{}^+ |\vec{N}(2)\rangle_V = \delta^{l2} \prod_{i \leq j}^2 (l'^{+}_{ij})^{n_{ij}} l'_2{}^+ |\vec{0}_{ij}, \vec{n}_s\rangle_V + \delta^{l1} m_1 |\vec{n}_{ij}, n_1 + 1, p_{12}, n_2\rangle_V, \quad l = 1, 2, \quad (47)$$

$$\begin{aligned} t'^{+}_{12} |\vec{N}(2)\rangle_V &= \prod_{i \leq j}^2 (l'^{+}_{ij})^{n_{ij}} t'^{+}_{12} |\vec{0}_{ij}, \vec{n}_s\rangle_V - 2n_{11} |n_{11} - 1, n_{12} + 1, n_{22}, \vec{n}_s\rangle_V \\ &\quad - n_{12} |n_{11}, n_{12} - 1, n_{22} + 1, \vec{n}_s\rangle_V. \end{aligned} \quad (48)$$

Here we have used the notations, first, $\vec{n}_s \equiv (n_1, p_{12}, n_2)$, $\vec{0}_{ij} \equiv (0, 0, 0)$ in accordance with definition of $|\vec{N}(2)\rangle_V$, second, $\delta_{ij,lm} = \delta_{il}\delta_{jm}$, for $i \leq j, l \leq m$, so that the vector $|\vec{N} + \delta_{ij,lm}\rangle_V$ in the Eq.(46) means subject to definition (44) increasing of only the coordinate n_{ij} in the vector $|\vec{N}(2)\rangle_V$, for $i = l, j = m$, on unit with unchanged values of the rest ones.

In turn, the action of Cartan-like generators on the vector $|\vec{N}(2)\rangle_V$ are given by the relations,

$$g'_0{}^l |\vec{N}(2)\rangle_V = (2n_l + n_{12} + n_l + (-1)^l p_{12} + h^l) |\vec{N}(2)\rangle_V, \quad l = 1, 2, \quad (49)$$

$$l'_0 |\vec{N}(2)\rangle_V = \prod_{i \leq j}^2 (l'^{+}_{ij})^{n_{ij}} l'_0 |\vec{0}_{ij}, \vec{n}_s\rangle_V. \quad (50)$$

To derive the Eqs. (46)–(50) we have used the formula for the product of operators A, B ,

$$AB^n = \sum_{k=0}^n C_k^n B^{n-k} \text{ad}_B^k A, \quad \text{ad}_B^k A = [\dots[A, \overbrace{B}^{k \text{ times}}], \dots], \quad \text{for } n \geq 0, \quad C_k^n = \frac{n!}{k!(n-k)!}. \quad (51)$$

At last, the action of the positive root vectors from the subspace \mathcal{E}_2^+ on the vector $|\vec{N}(2)\rangle_V$ being based on the rule (51), reads as follows,

$$\begin{aligned} t'_{12} |\vec{N}(2)\rangle_V &= \prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} \left(\frac{l'_1}{m_1} \right)^{n_1} (t'_{12})^{p_{12}} t'_{12} |\vec{0}_{ij}, 0, 0, n_2\rangle_V - \sum_l l n_{l2} |\vec{n}_{ij} - \delta_{ij,l2} + \delta_{ij,l1}, \vec{n}_s\rangle_V \\ &\quad + p_{12}(h^1 - h^2 - n_2 - p_{12} + 1) |\vec{n}_{ij}, n_1, p_{12} - 1, n_2\rangle_V, \end{aligned} \quad (52)$$

$$\begin{aligned} l'^1 |\vec{N}(2)\rangle_V &= -m_1 n_{11} |\vec{n}_{ij} - \delta_{ij,11}, n_1 + 1, p_{12}, n_2\rangle_V \\ &\quad + \left\{ \prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} l'^1 - \frac{n_{12}}{2} \prod_{i \leq j}^2 (l'_{ij})^{n_{ij} - \delta_{ij,12}} l'_2 \right\} |\vec{0}_{ij}, \vec{n}_s\rangle_V, \end{aligned} \quad (53)$$

$$\begin{aligned} l'^2 |\vec{N}(2)\rangle_V &= -m_1 \frac{n_{12}}{2} |\vec{n}_{ij} - \delta_{ij,12}, n_1 + 1, p_{12}, n_2\rangle_V \\ &\quad + \left\{ \prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} l'^2 - n_{22} \prod_{i \leq j}^2 (l'_{ij})^{n_{ij} - \delta_{ij,22}} l'_2 \right\} |\vec{0}_{ij}, \vec{n}_s\rangle_V, \end{aligned}$$

$$\begin{aligned} l'^{11} |\vec{N}(2)\rangle_V &= n_{11}(n_{11} + n_{12} + n_1 - p_{12} - 1 + h^1) |\vec{n}_{ij} - \delta_{ij,11}, \vec{n}_s\rangle_V \\ &\quad + \frac{n_{12}(n_{12} - 1)}{4} |\vec{n}_{ij} - 2\delta_{ij,12} + \delta_{ij,22}, \vec{n}_s\rangle_V \\ &\quad + \left\{ \prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} l'^{11} - \frac{n_{12}}{2} \prod_{i \leq j}^2 (l'_{ij})^{n_{ij} - \delta_{ij,12}} l'_{12} \right\} |\vec{0}_{ij}, \vec{n}_s\rangle_V, \end{aligned} \quad (54)$$

$$\begin{aligned} l'^{12} |\vec{N}(2)\rangle_V &= \frac{n_{12}}{4} \left(n_{12} + \sum_l (2n_{ll} + n_l + h^l) - 1 \right) |\vec{n}_{ij} - \delta_{ij,12}, \vec{n}_s\rangle_V \\ &\quad + \frac{1}{2} p_{12} n_{11} (h^2 - h^1 + n_2 + p_{12} - 1) |\vec{n}_{ij} - \delta_{ij,11}, n_1, p_{12} - 1, n_2\rangle_V \\ &\quad + n_{11} n_{22} |\vec{n}_{ij} - \delta_{ij,11} - \delta_{ij,22} + \delta_{ij,12}, \vec{n}_s\rangle_V \\ &\quad + \left\{ \prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} l'^{12} - \frac{n_{22}}{2} \prod_{i \leq j}^2 (l'_{ij})^{n_{ij} - \delta_{ij,22}} l'_{12} \right\} |\vec{0}_{ij}, \vec{n}_s\rangle_V \\ &\quad - \frac{n_{11}}{2} \prod_{i \leq j}^2 (l'_{ij})^{n_{ij} - \delta_{ij,11}} \left(\frac{l'_1}{m_1} \right)^{n_1} (t'_{12})^{p_{12}} t'_{12} |\vec{0}_{ij}, 0, 0, n_2\rangle_V \end{aligned} \quad (55)$$

$$\begin{aligned} l'^{22} |\vec{N}(2)\rangle_V &= n_{22}(n_{12} + n_{22} + p_{12} + n_2 - 1 + h^2) |\vec{n}_{ij} - \delta_{ij,22}, \vec{n}_s\rangle_V \\ &\quad + \frac{n_{12} p_{12}}{2} (p_{12} - 1 + h^2 - h^1 + n_2) |\vec{n}_{ij} - \delta_{ij,12}, n_1, p_{12} - 1, n_2\rangle_V \\ &\quad + \frac{n_{12}(n_{12} - 1)}{4} |\vec{n}_{ij} + \delta_{ij,11} - 2\delta_{ij,12}, \vec{n}_s\rangle_V \\ &\quad + \prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} \left(\frac{l'_1}{m_1} \right)^{n_1} l'^{22} |\vec{0}_{ij}, 0, p_{12}, n_2\rangle_V \\ &\quad - \frac{n_{12}}{2} \prod_{i \leq j}^2 (l'_{ij})^{n_{ij} - \delta_{ij,12}} \left(\frac{l'_1}{m_1} \right)^{n_1} (t'_{12})^{p_{12}} t'_{12} |\vec{0}_{ij}, 0, 0, n_2\rangle_V. \end{aligned} \quad (56)$$

It is easy to see that to complete the calculations in Eqs. (47), (48), (53)–(56) we need to find the action of positive root vectors l^m, l'^m , $m = 1, 2$, Cartan-like vector l'_0 , negative root vectors l_2^+, t_{12}^+ on the vector $|\vec{0}_{ij}, \vec{n}_s\rangle_V$, and the rest operators t_{12}', l'^{22} on an arbitrary vector $|\vec{0}_{lm}, 0, 0, n_2\rangle_V$ in terms of linear combinations of definite vectors. We solve this rather non-trivial technical problem explicitly with introducing auxiliary quantities which we call *primary block-operator* \hat{t}_{12}' given in (68) and *derived block-operators* $\hat{t}_{12}^+, \hat{l}_2^+, \hat{l}'_0, \hat{l}'_m, \hat{l}'_{m2}$, $m = 1, 2$ below whose concrete expressions will be shown with details of Verma module construction for the algebra under consideration in [47]. Thus, we may to formulate the result in the form of

Theorem 1. The Verma module for the non-linear second order algebra $\mathcal{A}'(Y(2), AdS_d)$ exists, is determined by the relations (46), (49), (57)–(66), expressed with help of primary \hat{t}_{12}' and derived block-operators $\hat{t}_{12}^+, \hat{l}_2^+, \hat{l}'_0, \hat{l}'_m, \hat{l}'_{m2}$, $m = 1, 2$ and has the final form,

$$t_{12}' |\vec{N}(2)\rangle_V = p_{12}(h^1 - h^2 - n_2 - p_{12} + 1) |\vec{n}_{ij}, n_1, p_{12} - 1, n_2\rangle_V - \sum_l l n_{l2} |\vec{n}_{ij} - \delta_{ij, l2} + \delta_{ij, l1}, \vec{n}_s\rangle_V + \hat{t}_{12}' |\vec{N}(2)\rangle_V, \quad (57)$$

$$t_{12}^+ |\vec{N}(2)\rangle_V = - \sum_l (3 - l) n_{l1} |\vec{n}_{ij} - \delta_{ij, l1} + \delta_{ij, l2}, \vec{n}_s\rangle_V + \hat{t}_{12}^+ |\vec{N}(2)\rangle_V, \quad (58)$$

$$l_l^+ |\vec{N}(2)\rangle_V = \delta^{l1} m_1 |\vec{N}(2) + \delta_{s,1}\rangle_V + \delta^{l2} \hat{l}_2^+ |\vec{N}(2)\rangle_V, \quad (59)$$

$$l'_0 |\vec{N}(2)\rangle_V = l'_0 |\vec{0}_{ij}, \vec{n}_s\rangle_V |_{[\vec{0}_{ij} \rightarrow \vec{n}_{ij}]}, \quad (60)$$

$$l'_1 |\vec{N}(2)\rangle_V = l'_1 |\vec{0}_{ij}, \vec{n}_s\rangle_V |_{[\vec{0}_{ij} \rightarrow \vec{n}_{ij}]} - m_1 n_{11} |\vec{n}_{ij} - \delta_{ij, 11}, \vec{n}_s + \delta_{s,1}\rangle_V - \frac{n_{12}}{2} \hat{l}_2^+ |\vec{n}_{ij} - \delta_{ij, 12}, \vec{n}_s\rangle_V, \quad (61)$$

$$l'_2 |\vec{N}(2)\rangle_V = l'_2 |\vec{0}_{ij}, \vec{n}_s\rangle_V |_{[\vec{0}_{ij} \rightarrow \vec{n}_{ij}]} - m_1 \frac{n_{12}}{2} |\vec{n}_{ij} - \delta_{ij, 12}, \vec{n}_s + \delta_{s,1}\rangle_V \quad (62)$$

$$- n_{22} \hat{l}_2^+ |\vec{n}_{ij} - \delta_{ij, 22}, \vec{n}_s\rangle_V, \quad (63)$$

$$l'_{11} |\vec{N}(2)\rangle_V = n_{11}(n_{11} + n_{12} + n_1 - p_{12} - 1 + h^1) |\vec{n}_{ij} - \delta_{ij, 11}, \vec{n}_s\rangle_V + \frac{n_{12}(n_{12} - 1)}{4} |\vec{n}_{ij} - 2\delta_{ij, 12} + \delta_{ij, 22}, \vec{n}_s\rangle_V + l'_{11} |\vec{0}_{ij}, \vec{n}_s\rangle_V |_{[\vec{0}_{ij} \rightarrow \vec{n}_{ij}]} - \frac{n_{12}}{2} \hat{l}_{12}^+ |\vec{n}_{ij} - \delta_{ij, 12}, \vec{n}_s\rangle_V, \quad (64)$$

$$l'_{12} |\vec{N}(2)\rangle_V = \frac{n_{12}}{4} \left(n_{12} + \sum_l (2n_{l1} + n_l + h^l) - 1 \right) |\vec{n}_{ij} - \delta_{ij, 12}, \vec{n}_s\rangle_V + \frac{1}{2} p_{12} n_{11} (h^2 - h^1 + n_2 + p_{12} - 1) |\vec{n}_{ij} - \delta_{ij, 11}, n_1, p_{12} - 1, n_2\rangle_V + n_{11} n_{22} |\vec{n}_{ij} - \delta_{ij, 11} - \delta_{ij, 22} + \delta_{ij, 12}, \vec{n}_s\rangle_V + l'_{12} |\vec{0}_{ij}, \vec{n}_s\rangle_V |_{[\vec{0}_{ij} \rightarrow \vec{n}_{ij}]} - \frac{n_{22}}{2} \hat{l}_{12}^+ |\vec{n}_{ij} - \delta_{ij, 22}, \vec{n}_s\rangle_V - \frac{n_{11}}{2} \hat{l}_{12} |\vec{n}_{ij} - \delta_{ij, 11}, \vec{n}_s\rangle_V, \quad (65)$$

$$l'_{22} |\vec{N}(2)\rangle_V = n_{22}(n_{12} + p_{12} + n_2 + n_{22} - 1 + h^2) |\vec{n}_{ij} - \delta_{ij, 22}, \vec{n}_s\rangle_V + \frac{n_{12} p_{12}}{2} (p_{12} - 1 + h^2 - h^1 + n_2) |\vec{n}_{ij} - \delta_{ij, 12}, \vec{n}_s - \delta_{s,12}\rangle_V + \frac{n_{12}(n_{12} - 1)}{4} |\vec{n}_{ij} + \delta_{ij, 11} - 2\delta_{ij, 12}, \vec{n}_s\rangle_V + \hat{l}_{22} |\vec{N}(2)\rangle_V - \frac{n_{12}}{2} \hat{l}_{12} |\vec{n}_{ij} - \delta_{ij, 12}, \vec{n}_s\rangle_V. \quad (66)$$

In deriving these relations the rule was used

$$\prod_{i \leq j}^2 (l'_{ij})^{n_{ij}} (l'_0, l'_n, l'_{lm}) \left| \vec{0}_{ij}, \vec{n}_s \right\rangle_V \equiv (l'_0, l'_n, l'_{lm}) \left| \vec{0}_{ij}, \vec{n}_s \right\rangle_V \big|_{[\vec{0}_{ij} \rightarrow \vec{n}_{ij}]}, \quad l, m, n = 1, 2, \quad l \leq m, \quad (67)$$

when the multipliers $(l'_{ij})^{n_{ij}}$ act as only raising operators on vectors $(l'_0, l'_k, l'_{km}) \left| \vec{0}_{ij}, \vec{n}_s \right\rangle_V$. The *primary block operator* \hat{t}'_{12} (corresponding to non-Lie part of t'_{12}) is determined as follows

$$\begin{aligned} \hat{t}'_{12} \left| \vec{N}(2) \right\rangle_V &= \sum_{k=0}^{[(n_2-1)/2]} \left\{ \sum_{m=0}^{[n_2/2]} \sum_{l=0}^{[n_2/2-(^1m+1)]} \dots \right. \\ &\quad \left[n_2/2 - \sum_{i=1}^{k-1} (^im+i) - (k-1) \right] \left[n_2/2 - \sum_{i=1}^{k-1} (^im+i) - k - m \right] \\ &\quad \dots \sum_{m=0}^{k-1} \sum_{l=0}^{k-1} (-1)^k \left(\frac{-2r}{m_2^2} \right)^{\sum_{i=1}^k (^im+i)+k} \\ &\quad \times C_{2^1m+1}^{n_2} C_{2^1l+1}^{n_2-2^1m-1} \dots C_{2^km+1}^{n_2-2(\sum_{i=1}^{k-1} (^im+i)+k-1)} C_{2^kl+1}^{n_2-2(\sum_{i=1}^{k-1} (^im+i)+k-1)-2^km-1} \\ &\quad \times \left| \hat{A}_{\vec{n}_{ij} + [\sum_{i=1}^k (^im+i)+k] \delta_{i2} \delta_{j2}, n_1, n, n_2-2[\sum_{i=1}^k (^im+i)+k]} \right\rangle_V \left. \right\}, \quad (68) \end{aligned}$$

with the vector $\left| \hat{A}_{\vec{N}(2)} \right\rangle_V$ given as,

$$\begin{aligned} \left| \hat{A}_{\vec{N}(2)} \right\rangle_V &= n_2 p_{12} \left| \vec{N}(2) - \delta_{s,12} \right\rangle_V \\ &\quad - \frac{m_1}{m_2} \sum_{m=0}^{[n_2/2]} \left(\frac{-2r}{m_2^2} \right)^{^1m} C_{2^1m+1}^{n_2} \left| \vec{N}(2) + ^1m \delta_{ij,22} + \delta_{s,1} - (2^1m+1) \delta_{s,2} \right\rangle_V \\ &\quad - \sum_{m=1}^{[n_2/2]} \left(\frac{-2r}{m_2^2} \right)^{^1m} \left\{ C_{2^1m}^{n_2} \left| \vec{N}(2) + \delta_{ij,11} + (^1m-1) \delta_{ij,22} + \delta_{s,12} - 2^1m \delta_{s,2} \right\rangle_V \right. \\ &\quad - \left[C_{2^1m}^{n_2} (h^2 - h^1 + 2p_{12}) + C_{2^1m+1}^{n_2} \right] \left| \vec{N}(2) + \delta_{ij,12} + (^1m-1) \delta_{ij,22} - 2^1m \delta_{s,2} \right\rangle_V \\ &\quad \left. + p_{12} C_{2^1m}^{n_2} (h^2 - h^1 + p_{12} - 1) \left| \vec{N}(2) + ^1m \delta_{ij,22} - \delta_{s,12} - 2^1m \delta_{s,2} \right\rangle_V \right\} \\ &\quad - \sum_{m=0}^{[n_2/2]} \sum_{l=0}^{[n_2/2-(^1m+1)]} \left(\frac{-2r}{m_2^2} \right)^{^1m+^1l+1} C_{2^1m+1}^{n_2} \left\{ \left[C_{2^1l+1}^{n_2-2^1m-1} (h^2 - h^1 + 2p_{12} + n_2 \right. \right. \\ &\quad \left. \left. - 2(^1m+^1l+1)) - C_{2^1l+2}^{n_2-2^1m-1} \right] \left| \vec{N}(2) + \delta_{ij,12} + (^1m+^1l) \delta_{ij,22} - 2(^1m \right. \right. \\ &\quad \left. \left. + ^1l+1) \delta_{s,2} \right\rangle_V - p_{12} C_{2^1l+1}^{n_2-2^1m-1} (h^2 - h^1 + n_2 - 2(^1m+^1l+1) + p_{12} - 1) \right. \\ &\quad \times \left| \vec{N}(2) + (^1m+^1l+1) \delta_{ij,22} - \delta_{s,12} - 2(^1m+^1l+1) \delta_{s,2} \right\rangle_V \\ &\quad \left. - C_{2^1l+1}^{n_2-2^1m-1} \left| \vec{N}(2) + \delta_{ij,11} + (^1m+^1l) \delta_{ij,22} + \delta_{s,12} - 2(^1m+^1l+1) \delta_{s,2} \right\rangle_V \right. \\ &\quad \left. + \frac{m_1}{m_2} C_{2^1l+2}^{n_2-2^1m-1} \left| \vec{N}(2) + (^1m+^1l+1) \delta_{ij,22} + \delta_{s,1} - 2(^1m+^1l+\frac{3}{2}) \delta_{s,2} \right\rangle_V \right\}. \quad (69) \end{aligned}$$

The obtained result has obvious consequences, first, in case of reducing of $\mathcal{A}'(Y(2), AdS_d)$ to the quadratic algebra $\mathcal{A}'(Y(1), AdS_d)$, given in Ref. [10, 12] for the vanishing components

$n_{l2} = n_2 = p_{12} = 0, l = 1, 2$ of an arbitrary VM vector $|\vec{N}(1)\rangle_V \equiv |n_{11}, 0, 0, n_1, 0, 0\rangle_V$. Second, for reducing the AdS space to Minkowski $R^{1,d-1}$ space when the non-linear algebra $\mathcal{A}'(Y(2), AdS_d)$ for $r = 0$ turns to the Lie algebra $(T'^2 \oplus T'^{2*} \oplus l'_0)(0) \oplus sp(4)$ for $k = 2$ in the Eqs.(43). For former case we derive known Verma module [12] from results of Theorem 1 whereas for latter one we will get new Verma module realization (see [47]) for above Lie algebra being different from one for $sp(4)$ in [24] and in [29] for $k = 2$.

3.2. Fock space realization of $\mathcal{A}'(Y(2), AdS_d)$

In this section, we will find on a base of constructed Verma module the realization of $\mathcal{A}'(Y(2), AdS_d)$ in formal power series in degrees of creation and annihilation operators $(B_a, B_a^+) = ((b_i, b_{ij}, d_{12}), (b_i^+, b_{ij}^+, d_{12}^+))$ in \mathcal{H}' whose number coincides to ones of second-class constraints among \mathcal{O}'_I , i.e. with $\dim(\mathcal{E}_2^- \oplus \mathcal{E}_2^+)$. It is solved following the results of [49] and algorithms suggested in [50] initially elaborated for a simple Lie algebra, then enlarged to a non-linear quadratic algebra $\mathcal{A}(Y(1), AdS_d)$ Ref. [10]. To this end, we make use of the mapping for an arbitrary basis vector of the Verma module and one $|\vec{n}_{ij}, \vec{n}_s\rangle$ of new Fock space \mathcal{H}' ,

$$|\vec{n}_{ij}, \vec{n}_s\rangle_V \longleftrightarrow |\vec{n}_{ij}, \vec{n}_s\rangle = \prod_{i \leq j}^2 (b_{ij}^+)^{n_{ij}} \prod_{l=1}^2 (b_l^+)^{n_l} (d_{12}^+)^{p_{12}} |0\rangle, \quad (70)$$

for $b_{ij}|0\rangle = b_l|0\rangle = d_{12}|0\rangle = 0, \quad (|\vec{N}(2)\rangle_{[\vec{N}=0]} \equiv |0\rangle).$

Here, vector $|\vec{n}_{ij}, \vec{n}_s\rangle$, for non-negative integers n_{ij}, n_l, p_{12} are the basis vectors of a Fock space \mathcal{H}' generated by 6 pair of bosonic (B, B^+) operators, being the basis elements of the Heisenberg algebra A_6 , with the standard (only non-vanishing) commutation relations

$$[B_a, B_b^+] = \delta_{ab} \iff [b_{ij}, b_{lm}^+] = \delta_{il,jm}, \quad [b_i, b_m^+] = \delta_{im}, \quad [d_{12}, d_{12}^+] = 1. \quad (71)$$

Omitting the peculiarities of the correspondence among the Verma module vectors (especially one of $|\hat{A}_{\vec{N}(2)}\rangle_V$ for $\vec{N}(2) = (0, \dots, 0, n_2)$ (69) and Fock space \mathcal{H}' vector $|\vec{0}_{ij}, 0, 0, n_2\rangle$ (see [47] for details) we formulate our basic result as the

Theorem 2. The oscillator realization of the non-linear algebra $\mathcal{A}'(Y(2), AdS_d)$ over Heisenberg algebra A_6 exists in terms of formal power series in degrees of creation and annihilation operators, is given by the relations (72)–(75), (77), (78)–(83) and expressed with help of *primary block-operator* \hat{t}_{12} (74), and *derived block-operators*, $\hat{t}_{12}^+, \hat{l}_0^+, \hat{l}_m^+, \hat{l}_{m2}^+, m = 1, 2$ (76), (84)–(88) as follows.

First, for the trivial negative root vectors, we have

$$l_1^+ = m_1 b_1^+, \quad l_{ij}^+ = b_{ij}^+, \quad g_0^i = 2b_{ii}^+ b_{ii} + b_{12}^+ b_{12} + (-1)^i d_{12}^+ d_{12} + b_i^+ b_i + h^i. \quad (72)$$

Second, for the operator $t'_{12}(B, B^+)$ and *primary block-operator* $\hat{t}_{12}(B, B^+)$ we obtain,

$$\begin{aligned} t'_{12} &= (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} - b_{11}^+ b_{12} - 2b_{12}^+ b_{22} + \hat{t}_{12}, \\ \hat{t}_{12} &= \sum_{k=0} \left[\sum_{l=1} \sum_{m=0} \dots \sum_{k=0} \sum_{l=0} (-1)^k \left(\frac{-2r}{m_2^2} \right)^{\sum_{i=1}^k (i m + i l) + k} \prod_{i=1}^k \frac{1}{(2^i m + 1)!} \frac{1}{(2^i l + 1)!} \right. \\ &\quad \times (b_{22}^+)^{\sum_{i=1}^k (i m + i l) + k} \left\{ b_2^+ d_{12} b_2 - \frac{m_1}{m_2} \sum_{m=0} \left(\frac{-2r}{m_2^2} \right)^m \frac{(b_{22}^+)^m}{(2m + 1)!} b_1^+ b_2^{2m+1} \right. \\ &\quad \left. + \sum_{m=1} \left(\frac{-2r}{m_2^2} \right)^m (b_{22}^+)^{m-1} \left[b_{12}^+ \left\{ \frac{h^2 - h^1 + 2d_{12}^+ d_{12}}{(2m)!} + \frac{b_2^+ b_2}{(2m + 1)!} \right\} \right. \right. \\ &\quad \left. \left. - \frac{b_{11}^+ d_{12}^+}{(2m)!} - \frac{b_{22}^+}{(2m)!} (h^2 - h^1 + d_{12}^+ d_{12}) d_{12} \right] b_2^{2m} \right\} \end{aligned} \quad (73)$$

$$\begin{aligned}
& - \sum_{m=0} \sum_{l=0} \left(\frac{-2r}{m_2^2} \right)^{m+l+1} \frac{1}{(2m+1)!} (b_{22}^+)^{m+l} \left[b_{12}^+ \left\{ \frac{h^2 - h^1 + 2d_{12}^+ d_{12} + b_2^+ b_2}{(2l+1)!} \right. \right. \\
& - \left. \frac{b_2^+ b_2}{(2l+2)!} \right\} - \frac{b_{11}^+ d_{12}^+}{(2l+1)!} - \frac{b_{22}^+}{(2l+1)!} (h^2 - h^1 + b_2^+ b_2 + d_{12}^+ d_{12}) d_{12} \\
& \left. + \frac{m_1}{m_2} \frac{b_{22}^+}{(2l+2)!} b_1^+ b_2 \right] b_2^{2(m+l+1)} \left\} (b_2)^{2(\sum_{i=1}^k (i_{m+i} l) + k)} \right\}^9. \tag{74}
\end{aligned}$$

Third, for the operators $t_{12}^+, l_2'^+$ and derived block-operator \hat{t}_{12}^+ we have,

$$t_{12}^+ = -2b_{12}^+ b_{11} - b_{22}^+ b_{12} + \hat{t}_{12}^+, \tag{75}$$

$$\begin{aligned}
\hat{t}_{12}^+ &= \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^m (b_{11}^+)^m \left\{ \frac{d_{12}^+}{(2m)!} - \frac{m_2}{m_1} \frac{b_2^+ b_1}{(2m+1)!} \right\} b_1^{2m} \\
& - \sum_{m=1} \left(\frac{-2r}{m_1^2} \right)^m (b_{11}^+)^{m-1} \left\{ b_{12}^+ \left[\frac{(h^2 - h^1 + 2d_{12}^+ d_{12} + b_2^+ b_2)}{(2m)!} - \frac{b_1^+ b_1}{(2m+1)!} \right] \right. \\
& \left. + b_{22}^+ \frac{(h^1 - h^2 - d_{12}^+ d_{12} - b_2^+ b_2)}{(2m)!} d_{12} + \frac{b_{22}^+}{(2m)!} \hat{t}_{12} \right\} b_1^{2m}, \tag{76}
\end{aligned}$$

$$\begin{aligned}
l_2'^+ &= m_1 \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^{m+1} (b_{11}^+)^m \left\{ b_{12}^+ \left[\frac{(h^2 - h^1 + 2d_{12}^+ d_{12} + b_2^+ b_2)}{(2m+1)!} - \frac{b_1^+ b_1}{(2m+2)!} \right] \right. \\
& \left. + b_{22}^+ \frac{(h^1 - h^2 - d_{12}^+ d_{12} - b_2^+ b_2)}{(2m+1)!} d_{12} - b_{11}^+ \frac{d_{12}^+}{(2m+1)!} \right\} b_1^{2m+1} \\
& + m_2 \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^m \frac{(b_{11}^+)^m b_2^+}{(2m)!} b_1^{2m} + m_1 \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m b_{22}^+}{(2m+1)!} \hat{t}_{12} b_1^{2m+1}. \tag{77}
\end{aligned}$$

Fourth, for the Cartan-like vector l_0' the representation holds,

$$\begin{aligned}
l_0' &= \hat{l}_0 + \frac{m_1}{2} \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m}{(2m+1)!} \left\{ b_{12}^+ \hat{l}_2 + b_{11}^+ \hat{l}_1 \right. \\
& - \left. \frac{m_2}{2} [\hat{t}_{12} + (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12}] b_2^+ \right\} b_1^{2m+1} \\
& - r \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^m (b_{11}^+)^m b_1^+ \left\{ \frac{2(h^1 - d_{12}^+ d_{12}) - 3}{(2m+1)!} + \frac{2b_1^+ b_1}{(2m+2)!} \right\} b_1^{2m+1} \\
& + \frac{1}{2} \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m}{(2m+2)!} \left\{ b_{11}^+ \hat{l}_0 - r b_{11}^+ ([h^1 - d_{12}^+ d_{12}][h^1 - d_{12}^+ d_{12} - 2] \right. \\
& - h^2 - d_{12}^+ d_{12} - b_2^+ b_2 + 2d_{12}^+ (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12}) \\
& - 2r b_{11}^+ \hat{t}_{12} d_{12}^+ + 8r b_{11}^+ b_{12}^+ \hat{l}_{12} + 4r (b_{12}^+)^2 \hat{l}_{22} + 2r b_{12}^+ [(h^1 - h^2 \\
& - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} + \hat{t}_{12}] \left. [h^2 + b_2^+ b_2 + d_{12}^+ d_{12} - 2] \right\} b_1^{2m+2} \\
& + \frac{r}{2} \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m}{(2m+2)!} b_{22}^+ \left\{ [(h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} + \hat{t}_{12}] \right.
\end{aligned}$$

⁹ one should be noted that in (74) for $k = 0$ there are no doubled sums and the products $\prod_{i=1}^0 \dots$ is equal to 1 and the only term $b_2^+ d_{12} b_2$ survive

$$\times (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} + t'_{12} \hat{t}'_{12} \Big\} b_1^{2m+2}. \quad (78)$$

Fifth, the operators $l'_m, l'_{lm}, l, m = 1, 2, l \leq m$, read as follows,

$$\begin{aligned} l'_1 = & -m_1 b_1^+ b_{11} - \frac{1}{2} l'^{2+} b_{12} + \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^m \frac{(b_{11}^+)^m}{(2m)!} \hat{l}_1 b_1^{2m} \\ & + \frac{1}{m_1} \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^m \frac{(b_{11}^+)^m}{(2m+1)!} \left\{ \hat{l}_0 - r \left[(h^1 - d_{12}^+ d_{12})(h^1 - d_{12}^+ d_{12} - 2) - h^2 \right. \right. \\ & \left. \left. - d_{12}^+ d_{12} - b_2^+ b_2 \right] - r d_{12}^+ \left[h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12} \right] d_{12} - r \hat{t}'_{12} d_{12}^+ \right\} b_1^{2m+1} \\ & - 4m_1 \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m b_{12}^+}{(2m+1)!} (4^m - \frac{1}{2}) \hat{l}_{12} b_1^{2m+1} \\ & + \frac{m_1}{2} \sum_{m=1} \left(\frac{-8r}{m_1^2} \right)^m (b_{11}^+)^{m-1} b_1^+ \left\{ \frac{1}{2} \frac{(2(h^1 - d_{12}^+ d_{12}) - 1)}{(2m)!} + \frac{b_1^+ b_1}{(2m+1)!} \right\} b_1^{2m} \\ & + \frac{1}{2} \sum_{m=1} \left(\frac{-2r}{m_1^2} \right)^m (4^m - 1) \frac{(b_{11}^+)^{m-1}}{(2m)!} \left\{ 2b_{12}^+ \hat{l}'^2 - m_1 b_1^+ \right. \\ & \left. - m_2 (\hat{t}'_{12} + h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} b_2^+ \right\} b_1^{2m} \\ & + \frac{m_1}{2} \sum_{m=1} \left(\frac{-2r}{m_1^2} \right)^{m+1} (4^m - 1) \frac{(b_{11}^+)^{m-1}}{(2m+1)!} \left\{ b_{11}^+ \hat{t}'_{12} d_{12}^+ - 4(b_{12}^+)^2 \hat{l}'_{22} \right. \\ & \left. - 2b_{12}^+ \left[(h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} + \hat{t}'_{12} \right] (h^2 + b_2^+ b_2 + d_{12}^+ d_{12} - 2) \right. \\ & \left. + b_{11}^+ d_{12}^+ (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} \right\} b_1^{2m+1} \\ & - \frac{m_1}{2} \sum_{m=1} \left(\frac{-2r}{m_1^2} \right)^{m+1} (4^m - 1) \frac{(b_{11}^+)^{m-1}}{(2m+1)!} b_{22}^+ \left\{ \left[(h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} \right. \right. \\ & \left. \left. + \hat{t}'_{12} \right] (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} + t'_{12} \hat{t}'_{12} \right\} b_1^{2m+1}, \end{aligned} \quad (79)$$

$$\begin{aligned} l'_2 = & \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^m \frac{(b_{11}^+)^m}{(2m+2)!} \hat{l}_2 (b_1^+)^2 b_1^{2m+2} - 2m_1 \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m}{(2m+1)!} \sum_k b_{1k}^+ \hat{l}_{k2} b_1^{2m+1} \\ & - 2m_1 \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m}{4} \hat{t}'_{12} \left\{ \frac{(h^1 + h^2 + b_2^+ b_2 - 2)}{(2m+1)!} + \frac{b_1^+ b_1}{(2m+2)!} \right\} b_1^{2m+1} \\ & - \frac{m_1}{2} \sum_{m=0} \left(\frac{-2r}{m_1^2} \right)^{m+1} (b_{11}^+)^m \left\{ \frac{(h^1 + h^2 + b_2^+ b_2 - 2)}{(2m+1)!} + \frac{b_1^+ b_1}{(2m+2)!} \right\} \times \\ & \times (h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) d_{12} b_1^{2m+1} - \frac{m_1}{2} b_1^+ b_{12} - l'^{2+}_{22} b_{22}, \end{aligned} \quad (80)$$

$$\begin{aligned} l'_{11} = & (b_{11}^+ b_{11} + b_{12}^+ b_{12} + b_1^+ b_1 - d_{12}^+ d_{12} + h^1) b_{11} + \frac{b_{22}^+}{4} b_{12}^2 - \frac{1}{2} \hat{t}'_{12} b_{12} \\ & + \frac{1}{2} \sum_{m=0} \left(\frac{-8r}{m_1^2} \right)^{m+1} \frac{(b_{11}^+)^m}{(2m+2)!} \left\{ \frac{1}{4r} \hat{l}_0 + b_{12}^+ \hat{l}_{12} - \frac{1}{4} \left[d_{12}^+ (2 - h^1 - h^2 - b_2^+ b_2) d_{12} \right. \right. \end{aligned}$$

$$\begin{aligned}
& +h^1(h^1-2)-h^2-b_2^+b_2+\hat{t}_{12}d_{12}^+]\Big\}b_1^{2m+2} \\
& -\frac{1}{2}\sum_{m=1}\left(\frac{-8r}{m_1^2}\right)^m(b_{11}^+)^{m-1}b_1^+\left\{\frac{h^1-d_{12}^+d_{12}-\frac{1}{2}}{(2m+1)!}+\frac{b_1^+b_1}{(2m+2)!}\right\}b_1^{2m+1} \\
& -\frac{1}{m_1}\sum_{m=0}\left\{\left(\frac{-8r}{m_1^2}\right)^m\frac{(b_{11}^+)^m}{(2m+1)!}\hat{l}_1'-\left(\frac{-2r}{m_1^2}\right)^m(4^m-1)\frac{(b_{11}^+)^{m-1}b_{12}^+}{(2m+1)!}\hat{l}_2'\right\}b_1^{2m+1} \\
& +\frac{1}{2}\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^m(4^m-1)(b_{11}^+)^{m-1}\frac{b_1^+b_1}{(2m+1)!}b_1^{2m} \\
& +\frac{m_2}{2m_1}\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^m(4^m-1)\frac{(b_{11}^+)^{m-1}}{(2m+1)!}\left\{t'_{12}+\sum_l lb_{1l}^+b_{l2}\right\}b_2^+b_1^{2m+1} \\
& -\frac{1}{2}\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^{m+1}(4^m-1)\frac{(b_{11}^+)^{m-1}}{(2m+2)!}\left[b_{11}^+d_{12}^+-2b_{12}^+(h^2+b_2^+b_2+d_{12}^+d_{12}-1)\right]\times \\
& \quad \times\left(h^1-h^2-d_{12}^+d_{12}-b_2^+b_2\right)d_{12}b_1^{2m+2} \\
& -\frac{1}{2}\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^{m+1}(4^m-1)\frac{(b_{11}^+)^{m-1}}{(2m+2)!}\left\{b_{11}^+\hat{t}_{12}d_{12}^+-4b_{11}^+b_{12}^+\hat{l}_{12}\right. \\
& \quad \left.-4(b_{12}^+)^2\hat{l}_{22}-2b_{12}^+\hat{t}_{12}(h^2+b_2^+b_2+d_{12}^+d_{12}-2)\right\}b_1^{2m+2} \\
& +\frac{1}{2}\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^{m+1}(4^m-1)\frac{(b_{11}^+)^{m-1}}{(2m+2)!}b_{22}^+\left\{\left[(h^1-h^2-b_2^+b_2-d_{12}^+d_{12})d_{12}+\hat{t}_{12}\right]\times\right. \\
& \quad \left.\times(h^1-h^2-b_2^+b_2-d_{12}^+d_{12})d_{12}+t'_{12}\hat{t}_{12}\right\}b_1^{2m+2}, \tag{81}
\end{aligned}$$

$$\begin{aligned}
l'_{12} &= \frac{1}{4}\left(b_{12}^+b_{12}+\sum_m(2b_{mm}^+b_{mm}+b_m^+b_m+h^m)\right)b_{12}+b_{12}^+b_{11}b_{22}-\frac{1}{2}\left(\hat{t}_{12}^+b_{22}+\hat{t}_{12}b_{11}\right) \\
& +\frac{1}{2}(h^2-h^1+b_2^+b_2+d_{12}^+d_{12})b_{11}d_{12}-\frac{1}{2m_1}\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^m\frac{(b_{11}^+)^m}{(2m+1)!}\hat{l}_2'b_1^{2m+1} \\
& +\sum_{m=0}\left(\frac{-2r}{m_1^2}\right)^m\frac{(b_{11}^+)^m}{(2m)!}\hat{l}_{12}b_1^{2m}+\sum_{m=1}\left(\frac{-2r}{m_1^2}\right)^m\frac{(b_{11}^+)^{m-1}b_{12}^+}{(2m)!}\hat{l}_{22}b_1^{2m} \\
& +\frac{1}{4}\sum_{m=1}\left(\frac{-2r}{m_1^2}\right)^m(b_{11}^+)^{m-1}\left\{\hat{t}_{12}\left[\frac{(h^1+h^2+b_2^+b_2-2)}{(2m)!}+\frac{b_1^+b_1}{(2m+1)!}\right]\right. \\
& \left.+\left[\frac{h^1+h^2+b_2^+b_2-2}{(2m)!}+\frac{b_1^+b_1}{(2m+1)!}\right](h^1-h^2-b_2^+b_2-d_{12}^+d_{12})d_{12}\right\}b_1^{2m}, \tag{82}
\end{aligned}$$

$$\begin{aligned}
l'_{22} &= \hat{l}_{22}+(b_{12}^+b_{12}+d_{12}^+d_{12}+b_2^+b_2+b_{22}^+b_{22}+h^2)b_{22}+\frac{1}{4}b_{11}^+b_{12}^2-\frac{1}{2}\hat{t}_{12}b_{12} \\
& +\frac{1}{2}(h^2-h^1+d_{12}^+d_{12}+b_2^+b_2)d_{12}b_{12}. \tag{83}
\end{aligned}$$

In the Eqs.(78)–(83), the *derived block-operators* $\hat{l}'_0, \hat{l}'_m, \hat{l}'_{m2}$, for $m = 1, 2$ are written as follows,¹⁰

$$\hat{l}'_0 = m_0^2 - r \sum_{m=0} \left(\frac{-8r}{m_1^2}\right)^m (b_{22}^+)^m \left\{ b_2^+ \left[\frac{1}{(2m+1)!} (2h^2 + 2d_{12}^+d_{12} - 1) \right. \right.$$

¹⁰ they are directly determined through their action on vector $|\vec{0}_{ij}, 0, p_{12}, n_2\rangle_V$, see [47] for details

$$\begin{aligned}
& + \frac{2b_2^+ b_2}{(2m+2)!} \Big] b_2 - 2 \frac{m_1}{m_2} \frac{b_1^+ d_{12}^+}{(2m+1)!} b_2 \Big\} b_2^{2m} \\
& + \frac{1}{2} \sum_{m=0} \left(\frac{-8r}{m_2^2} \right)^{m+1} \frac{(b_{22}^+)^m}{(2m+2)!} \Big\{ b_{22}^+ (m_0^2 - r[h^2(h^2-4) + h^1 - (d_{12}^+)^2 d_{12}^2 \\
& + 2d_{12}^+ d_{12}(h^1-2)]) + 2rb_{12}^+ d_{12}^+(h^1 - d_{12}^+ d_{12} - 2) + rb_{11}^+ (d_{12}^+)^2 \Big\} b_2^{2m+2} \\
& + 2r \sum_{m=0} \sum_{l=0} \left(\frac{-8r}{m_2^2} \right)^m \left(\frac{-2r}{m_2^2} \right)^{l+1} \frac{(b_{22}^+)^m}{(2m+1)!} \frac{(b_{22}^+)^{l+1}}{(2l+1)!} \hat{t}_{12}^+ d_{12}^+ b_2^{2(m+l+1)} \\
& + 2r \sum_{m=0} \sum_{l=0} \left(\frac{-8r}{m_2^2} \right)^m \left(\frac{-2r}{m_2^2} \right)^{l+1} \frac{(b_{22}^+)^{m+l}}{(2m+1)!} \Big\{ b_{12}^+ d_{12}^+ \left[\frac{[h^2 - h^1 + 2d_{12}^+ d_{12} + b_2^+ b_2 + 2]}{(2l+1)!} \right. \\
& - \frac{b_2^+ b_2}{(2l+2)!} \Big] - \frac{b_{11}^+ (d_{12}^+)^2}{(2l+1)!} + \frac{b_{22}^+ (d_{12}^+ d_{12} + 1)}{(2l+1)!} [h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}] \\
& \left. + \frac{m_1}{m_2} \frac{b_{22}^+ b_1^+ d_{12}^+}{(2l+2)!} b_2 \Big\} b_2^{2(m+l+1)}, \tag{84}
\end{aligned}$$

$$\hat{l}_1 = -\frac{m_2}{2} \sum_{m=1} \left(\frac{-2r}{m_2^2} \right)^{m+1} (b_{22}^+)^m d_{12}^+ \left\{ \frac{(h^1 + h^2 - 2)}{(2m+1)!} + \frac{b_2^+ b_2}{(2m+2)!} \right\} b_2^{2m+1}, \tag{85}$$

$$\begin{aligned}
\hat{l}_2 = & -\hat{l}_1 d_{12} - \sum_{m=0} \left(\frac{-8r}{m_2^2} \right)^m \frac{(b_{22}^+)^m}{(2m+1)!} \frac{[m_0^2 - rh^2(h^2-3)]}{m_2} b_2^{2m+1} \\
& + \frac{m_2}{4} \sum_{m=1} \left(\frac{-8r}{m_2^2} \right)^m (b_{22}^+)^{m-1} b_2^+ \left\{ \frac{(2h^2-1)}{(2m)!} + 2 \frac{b_2^+ b_2}{(2m+1)!} \right\} b_2^{2m} \\
& + \frac{m_2}{2} \sum_{m=1} \left(\frac{-2r}{m_2^2} \right)^{m+1} (b_{22}^+)^m d_{12}^+ \left\{ \frac{(h^1 + h^2 - 2)}{(2m+1)!} + \frac{b_2^+ b_2}{(2m+2)!} \right\} d_{12} b_2^{2m+1} \\
& - \frac{1}{2} \sum_{m=0} \left(\frac{-2r}{m_2^2} \right)^m (4^m - 1) (b_{22}^+)^{m-1} d_{12}^+ \left\{ \frac{m_1 b_1^+}{(2m)!} - \frac{m_2 b_2^+ d_{12}}{(2m)!} \right\} b_2^{2m} \\
& + \frac{m_2}{2} \sum_{m=0} \left(\frac{-2r}{m_2^2} \right)^{m+1} (4^m - 1) \frac{(b_{22}^+)^{m-1}}{(2m+1)!} \Big\{ b_{22}^+ [2(h^1-2)d_{12}^+ d_{12} \\
& + h^1 - h^2 - (d_{12}^+)^2 d_{12}^2] - 2b_{12}^+ d_{12}^+(h^1 - d_{12}^+ d_{12} - 2) - b_{11}^+ (d_{12}^+)^2 \Big\} b_2^{2m+1} \\
& - \frac{m_2}{2} \sum_{m=0} \sum_{l=0} \left(\frac{-2r}{m_2^2} \right)^{m+l+1} (4^m - 1) \frac{(b_{22}^+)^{m+l-1}}{(2m)!} \left\{ \frac{m_1}{m_2} \frac{b_{22}^+ b_1^+ d_{12}^+}{(2l+2)!} b_2 \right. \\
& + b_{12}^+ d_{12}^+ \left[\frac{(h^2 - h^1 + 2d_{12}^+ d_{12} + b_2^+ b_2 + 2)}{(2l+1)!} - \frac{b_2^+ b_2}{(2l+2)!} \right] \\
& \left. - \frac{(b_{11}^+ (d_{12}^+)^2 + b_{22}^+ [h^2 - h^1 + b_2^+ b_2])}{(2l+1)!} \right\} b_2^{2(m+l)+1} \\
& + \frac{m_2}{2} \sum_{m=0} \sum_{l=0} \left(\frac{-2r}{m_2^2} \right)^{m+l+1} (4^m - 1) \frac{(b_{22}^+)^m}{(2m)!} \frac{(b_{22}^+)^l}{(2l+1)!} \times \\
& \quad \times d_{12}^+ \left\{ h^2 - h^1 + b_2^+ b_2 + d_{12}^+ d_{12} + 2 \right\} d_{12} b_2^{2(m+l)+1} \\
& - \frac{m_2}{2} \sum_{m=0} \sum_{l=0} \left(\frac{-2r}{m_2^2} \right)^{m+l+1} (4^m - 1) \frac{(b_{22}^+)^m}{(2m)!} \frac{(b_{22}^+)^l}{(2l+1)!} \hat{t}_{12}^+ d_{12}^+ b_2^{2(m+l)+1}, \tag{86}
\end{aligned}$$

$$\hat{l}_{12} = \frac{1}{4} \sum_{m=1} \left(\frac{-2r}{m_2^2} \right)^m (b_{22}^+)^{m-1} d_{12}^+ \left\{ \frac{(h^1 + h^2 - 2)}{(2m)!} + \frac{b_2^+ b_2}{(2m+1)!} \right\} b_2^{2m}, \quad (87)$$

$$\begin{aligned} \hat{l}_{22} = & -2\hat{l}_{12}d_{12} - \sum_{m=0} \left(\frac{-8r}{m_2^2} \right)^m \frac{(b_{22}^+)^m}{m_2^2(2m+2)!} \{m_0^2 - rh^2(h^2 - 3)\} b_2^{2(m+1)} \\ & - \frac{1}{2} \sum_{m=1} \left(\frac{-8r}{m_2^2} \right)^m (b_{22}^+)^{m-1} b_2^+ \left\{ \frac{1}{2} \frac{(2h^2 - 1)}{(2m+1)!} + \frac{b_2^+ b_2}{(2m+2)!} \right\} b_2^{2m+1} \\ & + \frac{1}{2} \sum_{m=0} \left(\frac{-2r}{m_2^2} \right)^{m+1} (4^m - 1)(b_{22}^+)^{m-1} \left\{ \frac{b_{12}^+ d_{12}^+}{(2m+2)!} [2(h^1 - 2) - 2d_{12}^+ d_{12}] b_2 \right. \\ & + \frac{b_{22}^+}{(2m+2)!} [h^2 - h^1 - 2d_{12}^+ d_{12}(h^1 - 2) + (d_{12}^+)^2 d_{12}^2] b_2 + \frac{b_{11}^+ (d_{12}^+)^2 b_2}{(2m+2)!} \\ & \left. - \frac{b_2^+ d_{12}^+ d_{12}}{(2m+1)!} \right\} b_2^{2m+1} + \frac{m_1}{2m_2} \sum_{m=0} \left(\frac{-2r}{m_2^2} \right)^{m+1} \frac{(b_{22}^+)^{m-1} b_1^+ d_{12}^+}{(2m+1)!} (4^m - 1) b_2^{2m+1} \\ & + \frac{1}{2} \sum_{m=0} \sum_{l=0} \left(\frac{-2r}{m_2^2} \right)^{m+l+1} (4^m - 1) \frac{(b_{22}^+)^{m+l-1}}{(2m+1)!} \left\{ \frac{m_1}{m_2} \frac{b_{22}^+ b_1^+ d_{12}^+}{(2l+2)!} b_2 - \frac{b_{11}^+ (d_{12}^+)^2}{(2l+1)!} \right. \\ & + \frac{b_{22}^+}{(2l+1)!} (d_{12}^+ d_{12} + 1)(h^1 - h^2 - b_2^+ b_2 - d_{12}^+ d_{12}) \\ & \left. + b_{12}^+ d_{12}^+ \left[\frac{(h^2 - h^1 + 2d_{12}^+ d_{12} + b_2^+ b_2 + 2)}{(2l+1)!} - \frac{b_2^+ b_2}{(2l+2)!} \right] \right\} b_2^{2(m+l+1)} \\ & + \frac{1}{2} \sum_{m=0} \sum_{l=0} \left(\frac{-2r}{m_2^2} \right)^{m+l+1} (4^m - 1) \frac{(b_{22}^+)^{m+l}}{(2m+1)!(2l+1)!} \hat{t}_{12} d_{12}^+ b_2^{2(m+l+1)}. \end{aligned} \quad (88)$$

Note, the additional parts $o'_I(B, B^+)$ as the formal power series in the oscillators (B, B^+) do not obey the usual properties,

$$(l'_{lm})^+ \neq l'_{lm}, \quad (t'_{12})^+ \neq t'_{12}, \quad (l'_0)^+ \neq l'_0, \quad (l'_m)^+ \neq l'_m, \quad l \leq m, \quad (89)$$

if one should use the standard rules of Hermitian conjugation for the new creation and annihilation operators, $(B_a)^+ = B_a$. Restoration the proper Hermitian conjugation properties for o'_I , is achieved by changing the scalar product in \mathcal{H}' as follows,

$$\langle \Phi_1 | \Phi_2 \rangle_{\text{new}} = \langle \Phi_1 | K' | \Phi_2 \rangle, \quad (90)$$

for any vectors $|\Phi_1\rangle, |\Phi_2\rangle$ with some non-degenerate operator K' . This operator is determined by the condition that all the operators of the algebra must have the proper Hermitian properties with respect to the new scalar product,

$$\langle \Phi_1 | K' E^{-\prime\alpha} | \Phi_2 \rangle = \langle \Phi_2 | K' E^{\prime\alpha} | \Phi_1 \rangle^*, \quad \langle \Phi_1 | K' G' | \Phi_2 \rangle = \langle \Phi_2 | K' G' | \Phi_1 \rangle^*, \quad (91)$$

for $(E^{\prime\alpha}; E^{-\prime\alpha}) = (l'_{lm}, t'_{12}, l'_m; l_{lm}^+, t_{12}^+, l_m^+)$, $G' = (g_0^i, l'_0)$. The relations (91) lead to definition the operator K' , Hermitian with respect to the standard scalar product $\langle | \rangle$, in the form

$$K' = Z^+ Z, \quad Z = \sum_{(\vec{n}_{lm}, \vec{n}_s) = (\vec{0}, \vec{0})}^{\infty} |\vec{N}(2)\rangle_V \frac{1}{(\vec{n}_{lm})! (\vec{n}_s)!} \langle 0 | \prod_{r=1}^2 b_r^{n_r} d_{12}^{p_{12}} \prod_{l, m \geq l} b_{lm}^{n_{lm}}, \quad (92)$$

where $(\vec{n}_{lm})! = n_{11}! n_{12}! n_{22}!$, $(\vec{n}_s)! = n_1! n_2! p_{12}!$ and the normalization ${}_V \langle 0 | 0 \rangle_V = 1$ is supposed.

Table 3. The non-linear part of the converted algebra $\mathcal{A}_c(Y(2), AdS_d)$.

| $[\downarrow, \rightarrow]$ | L_0 | L^i | L^{i+} |
|-----------------------------|-----------------------------|------------------------------|--------------------------|
| L_0 | 0 | $-r\hat{\mathcal{K}}_1^{i+}$ | $r\hat{\mathcal{K}}_1^i$ |
| L^j | $r\hat{\mathcal{K}}_1^{j+}$ | \hat{W}_b^{ji} | \hat{X}_b^{ji} |
| L^{j+} | $-r\hat{\mathcal{K}}_1^j$ | $-\hat{X}_b^{ij}$ | $-\hat{W}_b^{ji+}$ |

The theorem 2 has the same consequences as ones from theorem 1 which concern, first, the flat limit of the algebra $\mathcal{A}'(Y(2), AdS_d)$ and therefore a new representation for Lie algebra $(T'^2 \oplus T'^{2*} \oplus l'_0)(0) \oplus sp(4)$. Second, modulo the oscillator pairs $b_{m2}, b_{m2}^+, b_2, b_2^+, d_{12}, d_{12}^+, m = 1, 2$, the obtained representation coincides with one for totally symmetric HS fields quadratic algebra $\mathcal{A}'(Y(1), AdS_d)$ in [10].

The set of the equations which compose the results of the Theorems 1, 2 represents the general solution of mentioned in the Introduction the second problem on the LF construction for mixed-symmetry HS tensors on AdS spaces with given mass and spin $\mathbf{s} = (s_1, s_2)$.

4. Construction of Lagrangian Actions

To construct Lagrangian formulation for the HS tensor field of fixed generalized spin $\mathbf{s} = (s_1, s_2)$ we should, initially, determine explicitly the composition law for the deformed algebra $\mathcal{A}_c(Y(2), AdS_d)$ (for arbitrary $k \geq 2$ see [47]), find BRST operator for the non-linear algebra $\mathcal{A}_c(Y(2), AdS_d)$ and, finally, reproduce properly gauge-invariant Lagrangian formulation for the basic bosonic field $\Phi_{(\mu)_{s_1}, (\nu)_{s_2}}$.

4.1. Explicit form for the algebra $\mathcal{A}_c(Y(2), AdS_d)$

To this end as in the case of the algebra $\mathcal{A}'(Y(2), AdS_d)$ of o'_I the only multiplication law for quadratic part of the initial algebra $\mathcal{A}(Y(k), AdS_d)$ for $k=2$ is changed, while its linear part is given by the same $sp(2k)$ algebra as for the maximal Lie subalgebra for $\mathcal{A}(Y(2), AdS_d)$ and $\mathcal{A}'(Y(2), AdS_d)$ with the same form of the commutators $[O_a, O_I]$ for $O_a \in sp(4)$. From the Eqs. (33), (34) and Table 1 the non-linear part of the algebra $\mathcal{A}_c(Y(2), AdS_d)$ can be restored in the form of the Table 3, Here the functions $\hat{\mathcal{K}}_1^i, \hat{W}_b^{ji}, \hat{X}_b^{ij}$ (therefore its Hermitian conjugated quantities $\hat{\mathcal{K}}_1^{i+}, \hat{W}_b^{ji+}$) are given as follows:

$$\hat{W}_b^{ij} = 2r\epsilon^{ij} \left\{ \sum_l (-1)^l [G_0^l - g_0^l] L^{12} - l'^{12} \sum_l (-1)^l G_0^l - [(T^{12} - t'^{12}) L^{11} - l'^{11} T^{12}] \right. \\ \left. + (T_{12}^+ - t_{12}^+) L^{22} - l'^{22} T_{12}^+ \right\}, \quad (93)$$

$$\hat{\mathcal{K}}_1^j = 4 \sum_{i=1}^2 \left\{ (L^{ji+} - l'^{ji+}) L^i - l'^i L^{ji+} \right\} + 2(L^{j+} - l'^{j+}) G_0^j - 2(g_0^j + \frac{1}{2}) L^{j+} \\ - 2 \left\{ [(L^{1+} - l'^{1+}) T_{12}^+ - t_{12}^+ L^{1+}] \delta^{j2} \delta^{i1} + [(L^{2+} - l'^{2+}) T^{12} - t'^{12} L^{2+}] \delta^{j1} \delta^{i2} \right\}, \quad (94)$$

$$\hat{X}_b^{ij} = \left\{ L_0 + r \left[\hat{K}_0^{0i} - 2g_0^i G_0^i + 4l'^{ii} L^{ii} + 4l'^{ii} L^{ii+} + \left(\hat{K}_0^{12} - t_{12}^+ T^{12} - t'^{12} T_{12}^+ \right. \right. \right. \\ \left. \left. \left. + 4l'^{12+} L^{12} + 4l'^{12} L^{12+} \right) \right] \right\} \delta^{ij} \\ - r \left\{ 4 \sum_l [(L^{jl+} - l'^{jl+}) L^{li} - l'^{li} L^{jl+}] + \left(\sum_l (G_0^{lj} - g_0^l) - 2 \right) T^{12} - t'^{12} \sum_l G_0^l \right\} \delta^{i2} \delta^{j1} \\ - r \left\{ 4 \sum_l [(L^{jl+} - l'^{jl+}) L^{li} - l'^{li} L^{jl+}] + \left(T_{ij}^+ - t_{ij}^+ \right) \sum_l G_0^l - \left(\sum_l g_0^l + 2 \right) T_{12}^+ \right\} \delta^{i1} \delta^{j2}. \quad (95)$$

The quantities $\hat{K}_0^{0i}, \hat{K}_0^{12}$ above are the same as ones in (26) for $k = 2$, but expressed in terms of O_I .

Following to our experience from study of the (super)algebra $\mathcal{A}_c(Y(1), AdS_d)$ in [10, 12, 13] when in order to find exact BRST operator, we choose the Weyl (symmetric) ordering for quadratic combinations of O_I in the r.h.s. of the Eqs. (93)–(95) as follows, $O_I O_J = \frac{1}{2}(O_I O_J + O_J O_I) + \frac{1}{2}[O_I, O_J]$. As the result, the Table 3 for such kind of ordering must contain the quantities $\hat{W}_{bW}^{ij}, \hat{K}_{1W}^j, \hat{X}_{bW}^{ij}$ (and $\hat{W}_{bW}^{ij+}, \hat{K}_{1W}^{j+}$) which read as

$$\hat{W}_{bW}^{ij} = r\epsilon^{ij} \left\{ \sum_l (-1)^l \mathcal{G}_0^l L^{12} + \mathcal{L}^{12} \sum_l (-1)^l \mathcal{G}_0^l - \mathcal{T}^{12} L^{11} - \mathcal{L}^{11} T^{12} + \mathcal{T}^{12+} L^{22} + \mathcal{L}^{22} T^{12+} \right\} \quad (96)$$

$$\begin{aligned} \hat{K}_{1W}^j = & \left\{ 2 \sum_l [\mathcal{L}^{jl+} L^l + \mathcal{L}^l L^{j1+}] + \mathcal{L}^{j+} G_0^j + \mathcal{G}_0^j L^{j+} - \{\mathcal{L}^{1+} T_{12}^+ + \mathcal{T}_{12}^+ L^{1+}\} \delta^{j2} \right. \\ & \left. - [\mathcal{L}^{2+} T^{12} + \mathcal{T}^{12} L^{2+}] \delta^{j1} \right\}, \end{aligned} \quad (97)$$

$$\begin{aligned} \hat{X}_{bW}^{ij} = & \left\{ L_0 + r \left[\mathcal{G}_0^i G_0^j - 2\mathcal{L}_{ii}^+ L^{ii} - 2\mathcal{L}^{ii} L_{ii}^+ + \frac{1}{2}(\mathcal{T}^{12} T_{12}^+ + \mathcal{T}_{12}^+ T^{12} - 4\mathcal{L}_{12}^+ L^{12} - 4\mathcal{L}^{12} L_{12}^+) \right] \right\} \delta^{ij} \\ & - r \left\{ 2 \sum_l^2 (\mathcal{L}^{1l+} L^{l2} + \mathcal{L}^{l2} L^{1l+}) + \frac{1}{2} \sum_l^2 \mathcal{G}_0^l T^{12} + \frac{1}{2} \mathcal{T}^{12} \sum_l^2 \mathcal{G}_0^l \right\} \delta^{i2} \delta^{j1} \\ & - r \left\{ 2 \sum_l^2 (\mathcal{L}^{l2+} L^{1l} + \mathcal{L}^{1l} L^{l2+}) + \frac{1}{2} \mathcal{T}_{12}^+ \sum_l^2 \mathcal{G}_0^l + \frac{1}{2} \sum_l^2 \mathcal{G}_0^l T_{12}^+ \right\} \delta^{i1} \delta^{j2}, \end{aligned} \quad (98)$$

with the notation \mathcal{O}_I for the quantity $\mathcal{O}_I = (O_I - 2o'_I)$. Note, the ordering quantities (96)–(98) do not contain linear terms (except for L_0 in r.h.s of the last relation) as compared to (93)–(95).

Thus, we derive the algebra of converted operators O_I underlying HS field subject to an arbitrary unitary irreducible AdS group representation in AdS space with spin $\mathbf{s} = (s_1, s_2)$ so that the problem now to find BRST operator for $\mathcal{A}_c(Y(2), AdS_d)$.

4.2. BRST-operator for converted algebra $\mathcal{A}_c(Y(2), AdS_d)$

The non-linear algebra now has not the form of closed algebra because of the operatorial functions $F_{IJ}^{(2)K}(o', O)$ in the Eq. (34) and as it was shown in [31] it leads to appearance of higher order structural functions due to the quadratic algebraic relations (96)–(98) and their Hermitian conjugates corresponding to those quantities $F_{IJ}^{(2)K}(o', O)$ for $i, j = 1, 2$. In ref.[31] it was found new structure functions $F_{IJK}^{RS}(O)$ of the 3rd order in terminology of Ref.[17], implied by a resolution of the Jacobi identities $[[O_I, O_J], O_K] + cycl.perm.(I, J, K) = 0$, as follows (32),

$$\left\{ F_{IJ}^M F_{MK}^P + [F_{IJ}^{(2)P}, O_K] + cycl.perm.(I, J, K) \right\} = F_{IJK}^{RS} \left(O_R \delta_S^P - \frac{1}{2} F_{RS}^P \right), \quad (99)$$

for $F_{IJ}^M \equiv (f_{IJ}^M + F_{IJ}^{(2)M})$. The structure functions $F_{IJK}^{RS}(o', O)$ are antisymmetric with respect to a permutation of any two of lower indices (I, J, K) and upper ones R, S and exist because of nontrivial Jacobi identities for the $k(2k - 1) = 6$ triples $(L_i, L_j, L_0), (L_i^+, L_j^+, L_0), (L_i, L_j^+, L_0)$.

The construction of a BFV-BRST operator Q' for $\mathcal{A}_c(Y(2), AdS_d)$ are considered in [31] and has the general form

$$Q' = \mathcal{C}^I [O_I + \frac{1}{2} \mathcal{C}^J (f_{JI}^P + F_{JI}^{(2)P}) \mathcal{P}_P + \frac{1}{12} \mathcal{C}^J \mathcal{C}^K F_{KJI}^{RP} \mathcal{P}_R \mathcal{P}_P]. \quad (100)$$

for the (\mathcal{CP}) -ordering for the ghost coordinates $\mathcal{C}^I = \{\eta_0, \eta_G^i, \eta_i^+, \eta_i, \eta_{ij}^+, \eta_{ij}, \vartheta_{12}, \vartheta_{12}^+\}$, and their

conjugated momenta $\mathcal{P}_I = \{\mathcal{P}_0, \mathcal{P}_G^i, \mathcal{P}_i, \mathcal{P}_i^+, \mathcal{P}_{ij}, \mathcal{P}_{ij}^+, \lambda_{12}^+, \lambda_{12}\}^{11}$. Explicitly, Q' given as,

$$\begin{aligned} Q' = & Q'_1 + Q'_2 + r^2 \left\{ \eta_0 \sum_{i,j} \eta_i \eta_j \varepsilon^{ij} \left[\frac{1}{2} \sum_m \left(G_0^m [\lambda_{12} \mathcal{P}_{22}^+ - \lambda_{12}^+ \mathcal{P}_{11}^+ + i \mathcal{P}_{12}^+ \sum_l (-1)^l \mathcal{P}_G^l] \right. \right. \right. \\ & \left. \left. - i(L_{11}^+ \lambda_{12}^+ - L_{22}^+ \lambda_{12}) \mathcal{P}_G^m + 4L^{mm} \mathcal{P}_{m2}^+ \mathcal{P}_{1m}^+ \right) - L_{12}^+ \mathcal{P}_G^1 \mathcal{P}_G^2 + 2L^{12} \mathcal{P}_{22}^+ \mathcal{P}_{11}^+ \right] \\ & + \eta_0 \sum_{i,j} \eta_i^+ \eta_j \left[\sum_m \left((-1)^m \frac{i}{2} G_0^m \sum_l \mathcal{P}_G^l + 2(L_{22}^+ \mathcal{P}^{22} - L^{11} \mathcal{P}_{11}^+) \right) \lambda_{12} \delta^{1j} \delta^{2i} \right. \\ & + \varepsilon^{\{1j\} \delta^{2i\}} \left(i \sum_m \left[\frac{1}{2} T^{12} \lambda_{12}^+ - 2L^{12} \mathcal{P}_{12}^+ (-1)^m \right] \mathcal{P}_G^m + 2 \left[L^{12} \lambda_{12} - T^{12} \mathcal{P}^{12} \right] \mathcal{P}_{22}^+ \right. \\ & \left. \left. + 2 \left[T_{12}^+ \mathcal{P}^{12} - L^{12} \lambda_{12}^+ \right] \mathcal{P}_{11}^+ \right) - T^{12} \left[\mathcal{P}_G^1 \mathcal{P}_G^2 \delta^{2i} \delta^{1j} + 2\mathcal{P}^{11} \mathcal{P}_{22}^+ \delta^{1i} \delta^{2j} \right] \right. \\ & \left. - 2 \sum_m (-1)^m \left[(G_0^m \mathcal{P}^{11} + i L^{11} \mathcal{P}_G^m) \delta^{1i} \delta^{2j} - (G_0^m \mathcal{P}^{22} + i L^{22} \mathcal{P}_G^m) \delta^{2i} \delta^{1j} \right] \mathcal{P}_{12}^+ \right] + h.c. \right\}, \end{aligned} \quad (101)$$

with the standard form for linear Q'_1 and quadratic Q'_2 terms in ghosts \mathcal{C}^I (see [31] for details). The Hermiticity of the nilpotent operator Q' in total Hilbert space $\mathcal{H}_{tot} = \mathcal{H} \otimes \mathcal{H}' \otimes \mathcal{H}_{gh}$ is defined by the rule,

$$Q'^+ K = K Q', \quad \text{for } K = \hat{1} \otimes K' \otimes \hat{1}_{gh}, \quad (102)$$

with the operator K' given in (92).

4.3. Lagrangian formulation

Properly the construction of Lagrangians for bosonic HS fields in AdS_d space, can be developed by partially following the algorithm of [11], [12], (see, as well [29]), which is a particular case of our construction, corresponding to $s_2 = 0$. As a first step, we extract the dependence of the BRST operator Q' (101) on the ghosts $\eta_G^i, \mathcal{P}_G^i$,

$$Q' = Q + \eta_G^i (\sigma^i + h^i) + \mathcal{B}^i \mathcal{P}_G^i \quad (103)$$

with some inessential in later operators \mathcal{B}^i and the BRST operator Q which corresponds only to the converted first-class constraints $\{O_I\} \setminus \{G_0^i\}$,

$$\begin{aligned} Q = & \frac{1}{2} \eta_0 L_0 + \sum_i \eta_i^+ L^i + \sum_{l \leq m} \eta_{lm}^+ L^{lm} + \vartheta_{12}^+ T^{12} + \frac{i}{2} \sum_l \eta_l^+ \eta^l \mathcal{P}_0 \\ & - \vartheta_{12}^+ \sum_n (1 + \delta_{1n}) \eta^{1n+} \mathcal{P}^{n2} + \vartheta_{12}^+ \sum_n (1 + \delta_{n2}) \eta^{n2} \mathcal{P}^{+1n} + \frac{1}{2} \sum_n \eta_{n2}^+ \eta^{1n} \lambda^{12} \\ & - \frac{1}{2} \sum_{l \leq m} (1 + \delta_{lm}) \eta^m \eta_{lm}^+ \mathcal{P}^l - [\vartheta_{12} \eta^{+2} + \vartheta_{12}^+ \eta^{+1}] \mathcal{P}^2 \\ & + r \left\{ \eta_0 \sum_i \eta_i^+ \left[2\mathcal{L}^{ii} \mathcal{P}_i^+ + 2\mathcal{L}^{i+} \mathcal{P}_{ii} + \mathcal{G}_0^i \mathcal{P}_i + 2(\mathcal{L}^{12} \mathcal{P}^{\{1+} + \mathcal{L}^{\{1+} \mathcal{P}_{12}\}} \delta^{2\} \}^i \right. \right. \\ & \left. \left. - \delta^{1i} (\mathcal{L}^2 \lambda_{12}^+ + \mathcal{T}_{12}^+ \mathcal{P}_2) - \delta^{2i} (\mathcal{L}^1 \lambda_{12} + \mathcal{T}_{12} \mathcal{P}_1) \right] \right. \\ & \left. - \frac{1}{2} \sum_{i,j} \eta_i^+ \eta_j^+ \varepsilon^{ij} \left[\sum_m (-1)^m \mathcal{G}_0^m \mathcal{P}_{12} - (\mathcal{T}_{12} \mathcal{P}_{11} + \mathcal{L}^{11} \lambda_{12}) + \mathcal{T}_{12}^+ \mathcal{P}_{22} + \mathcal{L}^{22} \lambda_{12}^+ \right] \right. \\ & \left. + 2\eta_i^+ \eta_j \left[\sum_m \mathcal{L}^{jm+} \mathcal{P}^{im} - \frac{1}{8} (\mathcal{T}_{12}^+ \lambda_{12} + \mathcal{T}_{12} \lambda_{12}^+) \delta^{ij} + \frac{1}{4} \sum_m \mathcal{G}_0^m \lambda_{12} \delta^{j1} \delta^{i2} \right] \right\} \\ & + r^2 \left\{ \eta_0 \sum_{i,j} \eta_i \eta_j \varepsilon^{ij} \left[\frac{1}{2} \sum_m \left(G_0^m [\lambda_{12} \mathcal{P}_{22}^+ - \lambda_{12}^+ \mathcal{P}_{11}^+] + 4L^{mm} \mathcal{P}_{m2}^+ \mathcal{P}_{1m}^+ \right) + 2L^{12} \mathcal{P}_{22}^+ \mathcal{P}_{11}^+ \right] \right. \\ & \left. + \eta_0 \sum_{i,j} \eta_i^+ \eta_j \left[2 \sum_m (L_{22}^+ \mathcal{P}^{22} - L^{11} \mathcal{P}_{11}^+) \lambda_{12} \delta^{1j} \delta^{2i} - 2T^{12} \mathcal{P}^{11} \mathcal{P}_{22}^+ \delta^{1i} \delta^{2j} \right] \right\} \end{aligned}$$

¹¹ they obey to independent nonvanishing anticommutation relations $\{\vartheta_{12}, \lambda_{12}^+\} = 1, \{\eta_i, \mathcal{P}_j^+\} = \delta_{ij}, \{\eta_{lm}, \mathcal{P}_{ij}^+\} = \delta_{li} \delta_{jm}, \{\eta_0, \mathcal{P}_0\} = i, \{\eta_G^i, \mathcal{P}_G^j\} = i \delta^{ij}$, possess the standard ghost number distribution, $gh(\mathcal{C}^I) = -gh(\mathcal{P}_I) = 1$, providing the property $gh(Q') = 1$, and have the Hermitian conjugation properties of zero-mode pairs, $(\eta_0, \eta_G^i, \mathcal{P}_0, \mathcal{P}_G^i)^+ = (\eta_0, \eta_G^i, -\mathcal{P}_0, -\mathcal{P}_G^i)$

$$\begin{aligned}
& +2\varepsilon^{\{1j\delta^2\}i} \left((L^{12}\lambda_{12} - T^{12}\mathcal{P}^{12})\mathcal{P}_{22}^+ + (T_{12}^+\mathcal{P}^{12} - L^{12}\lambda_{12}^+)\mathcal{P}_{11}^+ \right) \\
& -2\sum_m (-1)^m \left(G_0^m \mathcal{P}^{11} \delta^{1i} \delta^{2j} - G_0^m \mathcal{P}^{22} \delta^{2i} \delta^{1j} \right) \mathcal{P}_{12}^+ \Big] \Big\} + h.c.^{12}. \quad (104)
\end{aligned}$$

The generalized spin operator $\vec{\sigma} = (\sigma^1, \sigma^2)$, extended by the ghost Wick-pair variables,

$$\sigma^i = G_0^i - h^i - \eta_i \mathcal{P}_i^+ + \eta_i^+ \mathcal{P}_i + \sum_m (1 + \delta_{im}) (\eta_{im}^+ \mathcal{P}^{im} - \eta_{im} \mathcal{P}_{im}^+) + [\vartheta_{12}^+ \lambda^{12} - \vartheta^{12} \lambda_{12}^+] (-1)^i, \quad (105)$$

commutes with Q , $[Q, \sigma^i] = 0$. We choose a representation for Hilbert space \mathcal{H}_{tot} coordinated with decomposition (103) such that the operators $(\eta_i, \eta_{ij}, \vartheta_{12}, \mathcal{P}_0, \mathcal{P}_i, \mathcal{P}_{ij}, \lambda_{12}, \mathcal{P}_G^i)$ annihilate vacuum vector $|0\rangle$, and suppose that the field vectors $|\chi\rangle$ as well as the gauge parameters $|\Lambda\rangle$ do not depend on ghosts η_G^i ,

$$\begin{aligned}
|\chi\rangle &= \sum_n \prod_l^2 (b_l^+)^{n_l} \prod_{i \leq j}^2 (b_{ij}^+)^{n_{ij}} (d_{12}^+)^{p_{12}} (\eta_0^+)^{n_{f0}} \prod_{i,j,l \leq m, n \leq o}^2 (\eta_i^+)^{n_{fi}} (\mathcal{P}_j^+)^{n_{pj}} (\eta_{lm}^+)^{n_{flm}} (\mathcal{P}_{no}^+)^{n_{pno}} \\
&\times (\vartheta_{12}^+)^{n_{f12}} (\lambda_{12}^+)^{n_{\lambda 12}} |\Phi(a_i^+)_{(n)_l(n)_{ij}p_{12}}^{n_{f0}(n)_{fi}(n)_{pj}(n)_{flm}(n)_{pno}(n)_{f12}(n)_{\lambda 12}}}\rangle. \quad (106)
\end{aligned}$$

The brackets $(n)_{fi}, (n)_{pj}, (n)_{pno}$ in the Eq.(106) means, for instance, for $(n)_{pno}$ the set of indices $(n_{p11}, n_{p12}, n_{p22})$. The sum above is taken over n_l, n_{ij}, p_{12} and running from 0 to infinity, and over the rest n 's from 0 to 1. The Hilbert space \mathcal{H}_{tot} is decomposed into a direct sum of Hilbert subspaces with definite ghost number: $\mathcal{H}_{tot} = \bigoplus_{k=-6}^6 \mathcal{H}_k$. Denote by $|\chi^k\rangle \in \mathcal{H}_{-k}$, the state (106) with the ghost number $-k$, i.e. $gh(|\chi^k\rangle) = -k$. Thus, the physical state having the ghost number zero is $|\chi^0\rangle$, the gauge parameters $|\Lambda\rangle$ having the ghost number -1 is $|\chi^1\rangle$ and so on. For vanishing of all auxiliary creation operators B^+ and ghost variables $\eta_0, \eta_i^+, \mathcal{P}_i^+, \dots$ the vector $|\chi^0\rangle$ must contain only physical string-like vector $|\Phi\rangle = |\Phi(a_i^+)_{(0)_l(0)_{ij}0_{12}}^{(0)_{f0}(0)_{fi}(0)_{pj}(0)_{flm}(0)_{pno}(0)_{f12}(0)_{\lambda 12}}}\rangle$,

$$|\chi^0\rangle = |\Phi\rangle + |\Phi_A\rangle, \quad |\Phi_A\rangle \Big|_{[B^+=\eta_0=\eta_i^+=\mathcal{P}_i^+=\eta_{lm}^+=\mathcal{P}_{no}^+=\vartheta_{12}^+=\lambda_{12}^+=0]} = 0 \quad (107)$$

One can show, using the part of equations of motion and gauge transformations, that the vector $|\Phi_A\rangle$ can be completely removed (see Ref.[47]).

The equation for the physical state $Q'|\chi^0\rangle = 0$ and the tower of the reducible gauge transformations, $\delta|\chi\rangle = Q'|\chi^1\rangle, \delta|\chi^1\rangle = Q'|\chi^2\rangle, \dots, \delta|\chi^{(s-1)}\rangle = Q'|\chi^{(s)}\rangle$, lead to relations:

$$Q|\chi\rangle = 0, \quad (\sigma^i + h^i)|\chi\rangle = 0, \quad (\varepsilon, gh)(|\chi\rangle) = (0, 0), \quad (108)$$

$$\delta|\chi\rangle = Q|\chi^1\rangle, \quad (\sigma^i + h^i)|\chi^1\rangle = 0, \quad (\varepsilon, gh)(|\chi^1\rangle) = (1, -1), \quad (109)$$

$$\begin{aligned}
& \dots \quad \dots \quad \dots \\
\delta|\chi^{s-1}\rangle &= Q|\chi^s\rangle, \quad (\sigma^i + h^i)|\chi^s\rangle = 0, \quad (\varepsilon, gh)(|\chi^s\rangle) = (s \bmod 2, -s). \quad (110)
\end{aligned}$$

Here ε means for Grassmann parity and $s = 6$ is the maximal stage of reducibility for the massive bosonic HS field, because of subspaces $H_k = \emptyset$, for all integer $k \leq -7$. The middle set of equations in (108)–(110) determines the possible values of the parameters h^i and the eigenvectors of the operators σ^i . Solving spectral problem, we obtain a set of eigenvectors, $|\chi^0\rangle_{(n)_2}, |\chi^1\rangle_{(n)_2}, \dots, |\chi^s\rangle_{(n)_2}, n_1 \geq n_2 \geq 0$, and a set of eigenvalues,

$$\sigma_i |\chi\rangle_{(n)_k} = \left(n^i + \frac{d-1-4i}{2} \right) |\chi\rangle_{(n)_k}, \quad -h^i = n_i + \frac{d-1-4i}{2}, \quad i = 1, 2, \quad n_1 \in \mathbf{Z}, n_2 \in \mathbf{N}_0. \quad (111)$$

¹² here, in writing the coefficients depending on o'_I, O_I we have used the convention from Eqs.(96)–(98), for \mathcal{O}_I

It is easy to see that in order to construct Lagrangian for the field corresponding to a definite Young tableau (1) the numbers n_i must be equal to the numbers of the boxes in the i -th row of the corresponding Young tableau, i.e. $n_i = s_i$. Thus, the state $|\chi\rangle_{(s)_2}$ contains the physical field (7) and all its auxiliary fields. We fix some values of $n_i = s_i$. After substitution: $h^i \rightarrow h^i(s_i)$ operator $Q_{(s_1, s_2)} \equiv Q_{|h^i \rightarrow h^i(s_i)}$, is nilpotent on each subspace $H_{tot(s_1, s_2)}$ whose vectors satisfy to the Eqs.(108) for (111). Hence, the Lagrangian equations of motion (one to one correspond to Eqs.(2)–(5) for $k = 2$), a sequence of reducible gauge transformations have the form

$$Q_{(s_1, s_2)}|\chi^0\rangle_{(s_1, s_2)} = 0, \quad \delta|\chi^s\rangle_{(s_1, s_2)} = Q_{(s_1, s_2)}|\chi^{s+1}\rangle_{(s_1, s_2)}, \quad s = 0, \dots, 5. \quad (112)$$

Analogously to totally symmetric bosonic HS fields [11], [12] one can show that Lagrangian action for fixed spin $(n)_2 = (s)_2$ is defined up to an overall factor as follows

$$\mathcal{S}_{(s_1, s_2)} = \int d\eta_0 \langle \chi^0 | K_{(s_1, s_2)} Q_{(s_1, s_2)} | \chi^0 \rangle_{(s_1, s_2)}, \quad \text{for } |\chi^0\rangle \equiv |\chi\rangle. \quad (113)$$

where the standard scalar product for the creation and annihilation operators is assumed with measure $d^d x \sqrt{|g|}$ over AdS space. The vector $|\chi^0\rangle_{(s)_2}$ and the operator $K_{(s)_2}$ in (113) are respectively the vector $|\chi\rangle$ (106) subject to spin distribution relations (111) for HS tensor field $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}}(x)$ and operator K (102) where the substitution $h_i \rightarrow -(n_i + \frac{d-1-4i}{2})$ is done. The corresponding LF for bosonic field with spin s subject to $Y(s_1, s_2)$ is a reducible gauge theory of maximally $L = 6$ -th stage of reducibility.

One can prove that the equations of motion (112) indeed reproduces only the basic conditions (2)–(5) for HS fields with given spin (s_1, s_2) and mass. Therefore, the resulting equations of motion because of the representation (107) have the form,

$$L_0|\Phi\rangle_{(s)_2} = (l_0 + m_0^2)|\Phi\rangle_{(s)_2}, \quad (l_i, l_{ij}, t_{12})|\Phi\rangle_{(s)_2} = (0, 0, 0), \quad i \leq j. \quad (114)$$

The above relations permit one to determine the parameter m_0 in a unique way in terms of $h^i(s_i)$,

$$m_0^2 = m^2 + r \left\{ \beta(\beta + 1) + \frac{d(d-6)}{4} + \left(h^1 - \frac{1}{2} + 2\beta \right) \left(\left(h^1 - \frac{5}{2} \right) + \left(h^2 - \frac{9}{2} \right) \right) \right\}, \quad (115)$$

whereas the values of parameter m_1, m_2 remain by completely arbitrary and may be used to reach special properties of the Lagrangian for given HS field.

The general action (113) gives, in principle, a direct receipt to obtain the Lagrangian for any component field $\Phi_{(\mu^1)_{s_1}, (\mu^2)_{s_2}}(x)$ from general vector $|\chi^0\rangle_{(s)_2}$ since the only what we should do it is a computation of vacuum expectation values of products of some number of creation and annihilation operators.

5. Conclusion

In the paper we have derived the quadratic non-linear HS symmetry algebra for description of arbitrary integer HS fields on AdS-spaces with any dimensions and subject to k row Young tableaux $Y(s_1, \dots, s_k)$. It is shown the difference of the obtained algebras $\mathcal{A}(Y(k), AdS_d)$ for $k = 2$, $\mathcal{A}'(Y(2), AdS_d)$, $\mathcal{A}_c(Y(2), AdS_d)$ corresponding respectively to initial set of operators, their additional parts and converted set of operators within additive conversion procedure, is due to their pure non-linear parts, which are, in turn, connected to the AdS_d -radius $(\sqrt{r})^{-1}$ presence through the set of isometry AdS-space operators.

To obtain the algebras we start from an embedding of bosonic HS fields into vectors of an auxiliary Fock space, treat the fields as coordinates of Fock-space vectors and reformulate the theory in such terms. We realize the conditions that determine an irreducible AdS-group representation with a given mass and generalized spin in terms of differential operator constraints

imposed on the Fock space vectors. These constraints generate a closed non-linear algebra of HS symmetry, which contains, with the exception of k basis generators of its Cartan subalgebra, a system of first- and second-class constraints. Above algebra coincides modulo isometry group generators with its Howe dual $sp(2k)$ symplectic algebra. The construction of a correct Lagrangian description requires a deformation of the initial symmetry algebra, into algebra $\mathcal{A}_c(Y(2), AdS_d)$ introducing the algebra $\mathcal{A}'(Y(2), AdS_d)$.

We have generalized the method of construction of Verma module [8] from the case of Lie (super)algebras [49], [50], [29] and for quadratic algebra $\mathcal{A}'(Y(1), AdS_d)$ for totally-symmetric HS fields [10], [12] on to case of non-linear algebra underlying mixed-symmetric HS bosonic fields on AdS-space with two-row Young tableaux. The Theorem 1 presents our basic result in this relation. We show that as the byproduct of Verma module derivation the Poincare–Birkhoff–Witt theorem is valid in case of the algebra under consideration, therefore providing the lifting of the Verma module for Lie algebra $\mathcal{A}(Y(2), R^{1,d-1})$ [being isomorphic to $(T^2 \oplus T^{2*}) \rtimes sp(4)$] to one for quadratic algebra in a deformation parameter r . Of course, the same it is expected to be true for general algebra $\mathcal{A}'(Y(k), AdS_d)$, for which we suppose to obtain the explicit form of Verma module in the recursive procedure manner by means of new *primary and derived block-operators*, like $\hat{t}'_{12}, \hat{t}'_{12}^+$.

We have obtained the representation for the 15 generators of the algebra $\mathcal{A}'(Y(2), AdS_d)$ over Heisenberg–Weyl algebra A_6 as the formal power series in creation and annihilation operators, which in case of flat space limit ($r = 0$) takes the polynomial form, coinciding with earlier known results, at least for $m = 0$ [24] and appearing new one for massive case [51] and for $k = 2$ in [29]. The Theorem 2 finalizes our second basic result on solution of this Fock space realization problem for $\mathcal{A}'(Y(2), AdS_d)$ through Verma module construction approach.

On a base of BFV-BRST operator Q' which was found in Ref. [31] exactly up to third degree in ghost coordinates, for the nonlinear algebra $\mathcal{A}_c(Y(2), AdS_d)$ of 15 converted constraints O_I by analyzing the structure of Jacobi identities for them we present a proper construction of gauge-invariant Lagrangian formulations for the bosonic HS fields of given spin $\mathbf{s} = (s_1, s_2)$ and mass on AdS_d space. The corresponding Lagrangian formulation is at most 6-th stage reducible Abelian gauge theory and is given by the Eqs.(112),(113). The last relations may be considered as the final result in solution of the general problem to construct Lagrangian formulation for non-Lagrangian initial AdS-group irreducible representations relations which describe the bosonic HS field with two rows in Young tableaux. One should be noted the unconstrained Lagrangians for the free mixed-symmetry HS fields with two rows in Young tableaux on a AdS background have not been derived until now in both “metric-like” and “frame-like” formulations. These results to be seen as the first step to interacting theory, following in part to the research [52], [53].

Among the directions for application of the obtained results we point out the developing of the unconstrained formulation with minimal number of auxiliary fields for the basic HS field with two and more rows in the Young tableaux analogously to formulation given in [54], [55] for totally symmetric fields, which as well may be derived from the universal Lagrangian formulation suggested in the paper.

From a mathematical point of view the construction of the Verma module for the algebra $\mathcal{A}'(Y(2), AdS_d)$ open the possibility to study both its structure and search singular, subsingular vectors in it, so that it, in principle, will then permit to construct new (non-scalar) infinite-dimensional representations for given algebra. Besides, the above results permit to definitely understand the problems of (generalized) Verma module construction for HS symmetry algebras and superalgebras underlying HS bosonic and respectively fermionic fields on AdS-spaces subject to multi-row Young tableaux.

- [6] Bekaert X, Cnockaert S, Iazeolla C and Vasiliev M A Nonlinear higher spin theories in various dimensions *Preprint* hep-th/0503128
- [7] Fotopoulos A and Tsulaia M 2008 Gauge invariant lagrangians for free and interacting higher spin fields. A review of BRST formulation *Int.J.Mod.Phys. A* **24** 1–60 *Preprint* 0805.1346[hep-th]
- [8] Dixmier J 1974 *Algebres Enveloppantes* (Gauthier-Villars, Paris [on English: Dixmier J 1977 *Enveloping Algebras* (New York, North Holland)
- [9] Dobrev V K, 2007 Representations and characters of the Virasoro algebra and N=1 super-Virasoro algebras, *Preprint* 0709.0105[hep-th]
- [10] Burdick C, Navratil O and Pashnev A 2002 On the Fock space realizations of nonlinear algebras describing the high spin fields in AdS spaces *Preprint* hep-th/0206027
- [11] Buchbinder I L, Pashnev A and Tsulaia M 2001 Lagrangian formulation of the massless higher integer spin fields in the AdS background, *Phys. Lett. B* **523** 338–46, *Preprint* hep-th/0109067
- [12] Buchbinder I L, Krykhtin V A and Lavrov P M 2007 *Nucl. Phys. B* **762** 344–76 *Preprint* hep-th/0608005
- [13] Buchbinder I L, Krykhtin V and Reshetnyak A 2007 *Nucl. Phys. B* **787** 211–40 *Preprint* hep-th/0703049
- [14] Kuleshov A and Reshetnyak A 2009 Programming realization of symbolic computations for non-linear commutator superalgebras over the Heisenberg-Weyl superalgebra: data structures and processing methods *Preprint* 0905.2705[hep-th]
- [15] Fronsdal C 1979 *Phys. Rev. D* **20** 848–56
- [16] Fradkin E S and Vilkovisky G A 1975 *Phys. Lett. B* **55** 224–6; Batalin I A and Vilkovisky G A 1977 *Phys. Lett. B* **69** 309–12
- [17] Henneaux M 1985 *Phys. Rept.* **1** 126
- [18] Witten E 1986 *Nucl.Phys. B* **268** 253
- [19] Grigoriev M and Damgaard P H 2000 *Phys. Lett. B* **474** 323–30 *Preprint* hep-th/9911092
- [20] Gitman D M, Moshin P Yu and Reshetnyak A A 2005 Local Superfield Lagrangian BRST Quantization *J. Math. Phys.* **46** 072302-01–24 *Preprint* hep-th/0507160
- [21] Gitman D M, Moshin P Yu and Reshetnyak A A 2005 *Phys. Lett. B* **621** 295–308 *Preprint* hep-th/0507049
- [22] Faddeev L D and Shatashvili S L 1986 *Phys.Lett. B* **167** 225
- [23] Batalin I A and Tyutin I V 1991 *Int.J.Mod.Phys. B* **6** 3255
- [24] Burdick C, Pashnev A and Tsulaia M 2001 *Mod. Phys. Lett. A* **16** 731–46 *Preprint* hep-th/0101201
- [25] Buchbinder I L and Krykhtin V A 2005 *Nucl. Phys. B* **727** 537–63 *Preprint* hep-th/0505092
- [26] Buchbinder I L, Krykhtin V A and Pashnev A 2005 *Nucl. Phys. B* **711** 367–91 *Preprint* hep-th/0410215
- [27] Buchbinder I L, Krykhtin V A, Ryskina L L and Takata H 2006 *Phys. Lett. B* **641** 386–92
- [28] Moshin P Yu and Reshetnyak A A 2007 BRST approach to Lagrangian formulation for mixed-symmetry fermionic higher-spin fields *J. High Energy Phys.* JHEP10(2007)040 *Preprint* 0707.0386[hep-th]
- [29] Buchbinder I L and Reshetnyak A A 2011 General Lagrangian formulation for higher spin fields with arbitrary index symmetry. I. Bosonic fields *Preprint* 1110.5044[hep-th]
- [30] Howe R 1989 Transcending classical invariant theory *J. Amer. Math. Soc.* **3** 2
- [31] Reshetnyak A A 2008 Nonlinear Operator Superalgebras and BFV-BRST Operators for Lagrangian Description of Mixed-symmetry HS Fields in AdS Spaces *Preprint* 0812.2329[hep-th]
- [32] Buchbinder I L and Lavrov P M 2007 Classical BRST charge for nonlinear algebras *J.Math.Phys.* **48** 082306
- [33] Alkalaev K B 2001 *Phys. Lett. B* **519** 121–8; 2004 *Theor. Math. Phys.* **140** 1253–63
- [34] Alkalaev K B and Vasiliev M A 2003 N=1 supersymmetric theory of higher spin gauge fields in AdS(5) at the cubic level *Nucl. Phys. B* **655** 57–92 *Preprint* hep-th/0206068
- [35] Alkalaev K B, Shaynkman O V and Vasiliev M A 2004 On the frame-like formulation of mixed-symmetry massless fields in (A)dS(d) *Nucl. Phys. B* **692** 363–93 *Preprint* hep-th/0311164
- [36] Alkalaev K B, Shaynkman O V and Vasiliev M A 2005 Lagrangian formulation for free mixed-symmetry bosonic gauge fields in (A)dS(d) *J. High Energy Phys.* JHEP08(2005)069 *Preprint* hep-th/0501108
- [37] Zinoviev Yu 2009 Toward frame-like gauge invariant formulation for massive mixed symmetry bosonic fields *Nucl. Phys. B* **812** 46 *Preprint* 0809.3287[hep-th]
- [38] Zinoviev Yu 2010 Towards frame-like gauge invariant formulation for massive mixed symmetry bosonic fields. II. General Young tableau with two rows *Nucl.Phys. B* **826** 490–510 *Preprint* 0907.2140[hep-th].
- [39] Bastianelli F, Corradini O and Latini E 2008 Spinning particles and higher spin fields on (A)dS backgrounds *J. High Energy Phys.* JHEP11(2008)054 *Preprint* 0810.0188[hep-th]
- [40] Campoleoni A, Francia D, Mourad J and Sagnotti A 2009 *Nucl.Phys. B* **815** 289–357
- [41] Campoleoni A, Francia D, Mourad J and Sagnotti A 2010 *Nucl. Phys. B* **828** 405–514
- [42] Zinoviev Yu M 2010 Spin 3 cubic vertices in a frame-like formalism *J. High Energy Phys.* JHEP08(2010)084 *Preprint* 1007.0158[hep-th]
- [43] Boulanger N and Skvortsov E D 2011 *J. High Energy Phys.* JHEP09(2011)063 *Preprint* 1107.5028[hep-th]
- [44] Metsaev R R 2004 *Phys. Lett. B* **590** 95–104 *Preprint* hep-th/0312297

- [45] Buchbinder I L, Krykhtin V and Ryskina L 2009 *Mod. Phys. Lett. A* **24** 401–15 *Preprint* 0810.3467[hep-th]
- [46] Vasiliev M A 2004 *J. High Energy Phys.* JHEP12(2004)046, *Preprint* hep-th/0404124
- [47] C Burdik, A Reshetnyak, *Gauge-invariant Lagrangian Formulation for Mixed-Symmetry Higher Spin Bosonic Fields on AdS-spaces*, in preparation
- [48] Braverman A and Gaiitsgory D 1996 Poincare-Birkhoff-Witt theorem for quadratic algebras of Koszul type *J. Algebra* **181** 315-28
- [49] Burdik C, *Realizations of the real simple Lie algebras: the method of construction*, J. Phys. A: Math. Gen. 18 (1985) 3101–3112
- [50] Burdik C, Grozman P, Leites D and Sergeev A 2000 Realization of Lie Algebras and Superalgebras in Terms of Creation and Annihilation Operators: I, Theor. Math. Phys. **124**, 1048–58
- [51] Buchbinder I, Krykhtin V and Takata H 2007 *Phys. Lett. B* **656** 253–64
- [52] Buchbinder I L, Fotopoulos A, Petkou A and Tsulaia M 2006 *Phys. Rev. D* **74** 105018
- [53] Fotopoulos F, Irges N, Petkou A and Tsulaia M 2007 Higher Spin Gauge Fields Interacting with Scalar: The Lagrangian Cubic Vertex *J. High Energy Phys.* JHEP10(2007)021
- [54] Buchbinder I L, Galajinsky A V and Krykhtin V A 2007 *Nucl. Phys. B* **779** 155
- [55] Buchbinder I L and Galajinsky A V 2008 Quartet unconstrained formulation for massive higher spin fields *J. High Energy Phys.* JHEP11(2008)081, *Preprint* 0810.2852[hep-th]